## ARCTIC CLIMATE IMPACT ASSESSMENT

## **INDEX**

## **Preface**

The Arctic Climate Impact Assessment scientific report was published in 2005 by Cambridge University Press. The publication schedule and resources did not allow the inclusion of an index at that time. Given the length and complexity of this detailed, technical report, it was felt that an index would be a valuable reference tool.

In 2006, the ACIA Secretariat and the Cooperative Institute for Arctic Research (CIFAR) at the University of Alaska Fairbanks arranged for the production of this index. The index will not be published in print form but will be made available on the ACIA web site (<a href="www.acia.uaf.edu">www.acia.uaf.edu</a>) and the Arctic Monitoring and Assessment Programme (AMAP) web site (<a href="www.amap.no/acia">www.amap.no/acia</a>).

ACIA is an assessment of climate impacts, thus, index references to "projected impacts" or "projected change", unless otherwise stated, are references to "projected impacts of climate change" or "projected change as a result of climate change". In general, "projected change" refers to changes in physical climate variables, while "projected impacts" refers to the impacts of changes in various physical variables on humans and their activities and the ecosystems of the Arctic.

Place names, while appearing in the index, are not exhaustively indexed. Species are indexed under their common name when used in the report, with a cross-reference from the scientific name.

\_\_\_\_\_

Project Director: John E. Walsh, Director, Cooperative Institute for Arctic Research (CIFAR)

Project Manager: Patricia A. Anderson, Consultant

Indexer and Editor: Lelani Arris

A publication of: ACIA Secretariat and Cooperative Institute for Arctic Research University of Alaska Fairbanks June 2007

A	Aethia cristatella. see crested auklet
abalone (Haliotis rufuscens), 716	Aethia psittacula. see parakeet auklet
Abies sibirica. see Siberian fir	Aethia pusilla. see least auklet
Abies spp. see firs	afforestation, 798, 800
Abramis brama. see carp bream	agriculture, 806–814
abrupt climate change, 1018	in Alaska, 808
ACIA	in Canada, 807–808
conclusions, 1019–1020	climate limitations, 807–809
description, 2–10, 16–17	climate projections, approach, 809–810
recommendations for future assessments, 1018–1019	disease and insects, 813
regions, 2, 990, 1003 (see also Region 1; Region 2; Region 3;	in Finland, 808
Region 4)	global context, 806–807
actinarians, 483	and government policy, 813–814
actinomycetes, 540	growing degree-day analysis, 811–812
active layer. see permafrost	in Iceland, 809
adaptation to climate change, 949–950, 981, 1020	increases, impact on contaminants, 432
of agriculture, 806	management and adaptation, 805-806
by arctic residents, 609	in Norway, 808–809
by arctic species, 578–579	and predator control, 617
assessing, 9	present-day, 807-809
barriers to, 663–665	projected impacts, 188, 200, 807, 808–809, 810–814,
and co-management, 613	1002–1003, 1007, 1008, 1010
cultural, 1000	projected impacts on biodiversity, 554–555, 562–564
definition, 685, 951, 955	projected impacts on reindeer overgrazing, 601
by forest managers, 792	projected impacts on wildlife habitat, 1006
by freshwater fisheries, 415–416	research recommendations, 810, 854
of human-environment systems, 946–949	subsistence, 807
and human health, 865, 898–900	surprise responses, 804–806
by indigenous peoples, 558–559, 665–666, 955–957, 963, 993	in Sweden, 796–797
by Inuvialuit of Sachs Harbour, 668–670	and temperature, 786–787
by Saami, 81	vulnerability analysis, 957
by wildlife management systems, 599, 605, 617, 638, 640	Agrilus anxius. see bronze birch borer
by Yamal Nenets, 677	Ahnfeltia plicata, 481
adaptation to past climate variability and change	air freezing index, 910-912, 913-914, 933, 938-939. see also
by arctic species, 11	temperature, air
by boreal tree species, 788–789	air thawing index, 910–912, 913–914, 938–939. see also
by Dene people, 81	temperature, air
by indigenous peoples, 9, 15, 62, 64, 67, 90–91, 92, 651,	Aklavik, NWT, 654
662–663	akpiks. see cloudberry
by Inuit, 672	Alaria esculenta, 481
by Saami, 82, 89	Alaska
adaptive capacity, 980–981	agriculture, 808
of arctic species, 254–255, 266, 271, 284, 286, 542, 543, 548,	biodiversity, 1016
583	demographics, 868–869
of Atlantic herring, 485	forest area, 784
of boreal tree species, 553–554	forest tenure and management, 793–794
of commercial fisheries, 707, 708–709, 722, 729–730, 745,	observed change in glaciers and ice sheets, 204
768	projected impacts on boreal forest, 793–794
definition, 950	sanitation systems, 884
of Dene people, 80–81	spruce bark beetle, 831–833
of freshwater biota, 384–385, 386, 395, 398, 406, 433, 437	wildfire regimes and management, 838–839
of indigenous peoples, 76, 91, 92–93, 95, 658, 682–683,	wildlife management, 617–621, 632–634
685–686	Alaska and Inuvialuit Beluga Whale Committee, 612
of Inuit, 675	Alaska Eskimo Whaling Commission, 633, 637, 666
of Inuvialuit of Sachs Harbour, 670–671	Alaska Maritime National Wildlife Refuge, 604
of marine biota, 624	Alaska Native Claims Settlement Act, 618, 665, 793
of marine fish, 507, 544	Alaska plaice (Pleuronectes quadrituberculatus), 487, 751
of marine mammals, 509, 545, 624	albedo
of reindeer husbandry, 794–795	of boreal forest, 785, 789
research recommendations, 1019–1020	feedbacks, 998, 999
of Saami reindeer husbandry, 969	projected change, 1006
of Sachs Harbour, NWT, 963	projected impacts of change in snow cover, 200
of seabirds, 509, 545	and sea ice, 459, 474, 476–477, 519, 999, 1005
of social systems, 15, 729–730, 770–771	of terrestrial ecosystems, 314, 319
and vulnerability analysis, 957-958	of tundra, 550, 785
of wildlife management systems, 620	and UV radiation, 153, 160, 163-164, 166-167, 278, 995, 996
of Yamal Nenets, 676, 678	and vegetation change, 322, 324
adfreeze bond, 909, 914	Alca torda. see razorbill
aerosols. see also arctic haze	Alces alces. see moose
latitudinal gradient, 163	alcids. see seabirds
and ozone depletion, 155	alcohol abuse, 896, 897
and UV radiation, 153, 160, 162-163, 167, 174, 995	alder (Alnus spp.), 257, 271, 293, 371, 799

Aleut, 76–77, 633, 652, 681	Anser fabalis. see bean goose
Aleut International Association, 13, 68–71, 76–77, 602, 968	Anser fabalis fabialis. see taiga bean goose
Aleutian Islands, 605	Anser fabalis rossicus. see tundra bean goose
indigenous knowledge case study, 76–77	Antarctic Bottom Water, 473
Aleutian tern (Sterna aleutica), 635	apatite. see mineral extraction
alewife (Alosa spp.), 410	Apherusa glacialis, 480, 482
alfalfa, 807, 810, 812	aphids (Acyrthosiphon spp.), 268, 296
algae, 284, 549. see also microorganisms	aquaculture
adaptation to arctic environment, 270–271	in Bering Sea region, 747
diversity, 262	in Greenland, 716
impacts of oil pollution, 565	in Iceland, 716
knowledge of arctic species, 540	impacts on biodiversity, 555, 564, 576
projected change in distribution, 569	impacts on wild fish market, 765, 767
in sea ice, 480	in northeast Atlantic, 695
species numbers, 580	in northwest Atlantic, 739
alien species. <i>see</i> non-native species	in Norway, 702–703, 704, 706
Alle alle. see little auk	projected impacts, 417, 708, 745, 770
Allium schoenoprasum, 258	in Russia, 703
*	
Alnus spp. see alder	and wild species, 716, 880
Alopecosa hirtipes, 258	arachnids, 540, 580
Alopecurus alpinus. see alpine foxtail	Arctic, definition of, 2
Alopex lagopus. see Arctic fox	by CAFF, 259
Alosa spp. see alewife	fisheries regions, 693, 695, 710, 731, 746
alpine bistort (Polygonum viviparum), 247	for forests, 784
alpine foxtail (Alopecurus alpinus), 278	for freshwater ecosystems, 355
alpine saxifrage (Saxifraga nivalis), 247	marine Arctic, 26, 27, 454
alpine timothy (Phleum alpinum), 285	for model projections, 110
Alutiiq, 652, 681	for past and present climate, 22
Alvania sp., 483	in Russia, 624–625
American plaice. see long rough dab	for terrestrial ecosystems, 244–245
American robin (Turdus migratorius), 70–71, 260, 585	Arctic Athabaskan Council, 13, 68–71, 77–78, 79, 602, 968
American wigeon (Anas americana), 300	Arctic Borderlands Ecological Knowledge Co-op, 646, 684–685
Ammodytes spp. see sand lance	Arctic char (Salvelinus alpinus), 11, 570
ammonia, 312	adaptive capacity, 386
amphibians, 282, 571. see also wildlife	anadromous behavior, 409
amphipods, 480, 482, 483, 488, 489, 506	aquaculture, 417, 716
Anarhichas lupus lupus. see wolffish	biodiversity monitoring, 586
Anas acuta. see pintail	distribution, 396–397
Anas americana. see American wigeon	fisheries importance, 410
Anas crecca. see common teal	genetic diversity, 563–564
Anas penelope, see Eurasian wigeon	in Greenland, 388–389
Anas platyrhynchos, see mallard	harvest by indigenous peoples, 652, 654, 667, 672
anchovy (Engraulis spp.), 508, 703. see also northern anchovy	importance of estuaries, 361
ancient murrelet (Synthliboramphus antiquus), 633, 635	indigenous observations of change, 70–71, 83
Andromeda polifolia, 302	in Lake Saanajärvi, 382
anemia, 883	life history and environment, 400–401
Anguilla anguilla. see eel	paleoecology, 251
animals. see also birds; fish; mammals, marine; mammals,	projected change in distribution, 407, 409
terrestrial; individual species	projected impacts, 373, 436, 1010, 1012–1013, 1017
adaptations to arctic environment, 266–269	projected impacts on fisheries, 411–412, 417
community response to experimental warming, 296–297	temperature and metal accumulation, 429
diversity, present day, 259–260, 1016	and water temperature, 385, 398
genetic response to change, 285–286	Arctic cisco (Coregonus autumnalis), 361, 395, 410, 436
impacts of projected UV radiation change, 281–282	Arctic Coastal Dynamics (ACD) project, 219, 232, 921–922
impacts of recent climate change, 279–280	Arctic cod (Arctogadus glacialis), 70-71, 486, 495, 496, 758. see
indigenous observations of change, 70–71, 287, 660	also polar cod (Boreogadus saida)
predator-prey interactions, 301–302	Arctic Council, 6, 560, 623, 666, 686–687, 956
projected distribution change, 287–289, 290–291	Permanent Participants, 13, 602
projected impacts, 281	Arctic dryad (Dryas integrifolia), 249, 278
projected impacts on cyclic populations, 300–301	Arctic Environmental Protection Strategy (AEPS), 956
projected impacts on disease, 302–303	Arctic flounder ( <i>Pleuronectes glacialis</i> ), 385
projected impacts on parasites, 302–303	Arctic fox (Alopex lagopus), 619, 670
vulnerability, 269	adaptation to arctic environment, 266–267, 268
aniuvak. see snow beds	displacement by red fox, 281, 548
	· · · · · · · · · · · · · · · · · · ·
Anser albifrons, see white-fronted goose	genetic diversity, 285–286
Anser albifrons flavirostris. see Greenland white-fronted goose	harvest by indigenous peoples, 653, 667
Anser anser. see greylag goose	harvest in Russia, 607, 609
Anser brachyrhynchus. see pink-footed goose	impacts of industrial development, 677
Anser caerulescens. see snow goose	impacts of late freeze-up, 75, 76
Anser caerulescens caerulescens. see lesser snow goose	as invasive species, 605
Anser canagicus. see emperor goose	as mustelid predator, 300–301, 550
Anser erythropus. see lesser white-fronted goose	observed change, 608, 1005

projected impacts, 882, 1006	Arctogadus glacialis. see Arctic cod
as seabird predator, 489, 576, 605	Arctophila fulva, 258
and UV radiation, 268	Arenaria interpres. see ruddy turnstone
Arctic grayling (Thymallus arcticus), 373, 378, 388, 395, 402, 403, 407, 414, 436, 652	arrowtooth flounder (Atheresthes stomias), 487, 495, 502, 503, 503, 748, 755, 756, 758
Arctic hare (Lepus arcticus), 71, 266–267, 302, 654, 667	Artemisia spp., 293
arctic haze, 106, 143, 162–163, 429–430	Arthrobacter, 260–261, 283
Arctic lands, area, 245–246	arthropods, 552, 557
Arctic lemming, 258	Arviat, Nunavut, Canada, 68, 70
Arctic loon (Gavia arctica), 363, 633, 635	ascidians, 756
Arctic Mediterranean, 454, 461, 478	Asio flammeus. see short-eared owl
Arctic Monitoring and Assessment Programme (AMAP), 6, 652,	aspen (Populus tremuloides), 551-552, 792, 823, 833, 834, 836,
952, 959	845
Arctic National Wildlife Refuge (ANWR), 604, 621	aspen leaf miner (Phyllocnistis populiella), 834
Arctic Ocean. see also marine Arctic	Asplanchna priodonta, 422
area of, 26	assessment, science, 7–8
boundary, 26	Asteraceae, 293
and carbon sequestration, 34	Athapaskans, 652, 655–656, <i>681</i>
carbon storage, 1014	Atheresthes stomias, see arrowtooth flounder
change in "freshwater" storage, 45	Atka mackerel (Pleurogrammus monoptergius), 758, 760, 767
circulation, 27–30 climate model representation, 106–107	Atlantic cod (Gadus morhua), 486
described, 10	aquaculture, 745 around Iceland and Greenland, 710
"freshwater" defined, 27	in central North Atlantic, 711–712
freshwater inflows, 27–29	commercial fishery, 698–699, 704–707, 714, 715, 722–723,
influence of temperate latitudes, 27–29	736–737, 742–744, 765
observed change, 44–45, 923–924	distribution, 484
physical characteristics, 29–30	economic importance, 741
productivity, 493	export value, 704
projected impacts, 187–188, 188, 208, 506, 1008	historic fisheries, 695, 696–697, 710, 712–713, 732, 733
thresholds of change, 473–475	impacts of past climate variability, 501, 700, 716–717,
variability of temperature and salinity, 467	718–719, 736–737, 866
Arctic Ocean Model Intercomparison Project (AOMIP), 109, 135	impacts of temperature, 495, 700
Arctic Oscillation (AO), 10, 23, 24-25, 1018. see also North	impacts of UV radiation, 513, 515
Atlantic Oscillation	in northwest Atlantic, 732–733
and air temperature, 25, 39	overharvesting, 561, 700, 736–737
attribution of change, 25	past distribution change, 500
and Bering Sea fish, 502, 503	as predator of crab, 739
and contaminant transport, 890–891 and cyclone activity, 25	as predator of shrimp, 719, 738
impacts on ocean temperature, 572	as prey, 696 productivity, 484
and ocean circulation, 467	projected change in distribution, 545
projected change, 119, 127, 143, 470, 472–473	projected impacts, 507, 508, 700, 708, 721, 728, 740, 741,
and sea ice, 44, 192, 457–458, 471	1006–1007, 1013
and stratospheric ozone, 143	and seal predation, 739
and ungulates, 974	temperature and migration, 496
and winter precipitation, 279	Atlantic halibut (Hippoglossus hippoglossus), 652, 703, 716
Arctic poppy (Papaver polare), 556	Atlantic herring (Clupea harengus), 484-485
Arctic redpoll (Carduelis hornemanni), 267	in Central North Atlantic, 711
indigenous observations of change, 71	commercial fishery, 697–698, 705, 714, 722, 734
Arctic Regional Climate Model Intercomparison Project	distribution, 484
(ARCMIP), 109, 130, 131	export value, 704
arctic residents. see also indigenous peoples	historic fisheries, 696–697, 714
adaptation to climate change, 609, 613	impacts of past climate variability, 500, 500, 501, 700,
demographics, 13–14, 866, 867, 868 economic impacts, 1000, 1002	717–718, 738
economic impacts, 1000, 1002 ecosystem impacts, 247	impacts of temperature, 495 impacts of UV radiation, 514
impacts on ecosystems, 252, 545	in northeast Atlantic, 696
impacts on habitats, 554–555, 566–567	in northwest Atlantic, 732
multiple stresses, 946–949, 1000	Norwegian spring-spawning, 485, 496, 501, 698, 707
projected impacts, 15, 188, 195, 1000–1001, 1007, 1008–1009,	overharvesting, 561, 700
1010–1011, 1013–1014	as prey, 486, 488, 489, 572
research recommendations, 1019	projected change in distribution, 545
societal conditions and trends, 959	projected impacts, 507–508, 700, 708, 720, 741, 1006–1007
20th-century immigration, 13	Atlantic meridional overturning circulation. see thermohaline
Arctic skua. see parasitic jaeger	circulation
Arctic sorrel (Oxyria digyna), 271, 272	Atlantic puffin (Fratercula arctica), 489, 556, 572, 627, 629, 63
Arctic tern (Sterna paradisaea), 71, 489, 627, 629, 630, 631, 633,	Atlantic salmon (Salmo salar), 11, 359
635, 653	aquaculture, 417, 576, 702–703, 767
Arctic warbler (Phylloscopus borealis), 258	fisheries regulation, 417
Arctic willow (Salix arctica), 247, 258, 278, 294–295 Arctocetraria nigricascens, 258	harvest by indigenous peoples, 652 importance of estuaries, 361
Arcioceiraria nigricascens, 230	importance of estuaries, 301

life history, 544	commercial fisheries, 695–709
in northwest Atlantic, 732	ecosystem description, 696
past distribution change, 500	factors affecting ecosystem, 490, 544
projected impacts, 408–409, 1012	"Loophole," 695
projected range extensions, 407, 410	nutrient levels, 491
and water temperature, 411, 496	past climate variability and fish, 501
Atlantic saury (Scomberesox saurus), 500	productivity, 481, 483, 491, 492–493
atmosphere-ocean general circulation model (AOGCM), 102–109.	projected impacts, 122, 475–476, 505–506, 1005
see also climate models; climate models, ACIA-	projected impacts on fish distribution, 508
designated	variability of temperature and salinity, 467–468
atmospheric components of, 105–106	Barents Sea Impact Study (BASIS), 947
climate drift, 105	bark beetles, 834. see also spruce bark beetle; spruce engraver
coupling, 104–105	beetle
cryosphere component, 107–108	barley, 807–809, 810, 811, 852
ensemble simulations, 143–144	barnacle goose (Branta leucopsis), 288, 574, 575, 617, 630, 1005
flux corrections, 105	barnacles, 483
grid size, 105	Barrow, Alaska, 68 basking shark (Cetorhinus maximus), 500
improving projections, 140–144 initialization, 104–105	beach ice, 921
land-surface component, 107	bean goose (Anser fabalis), 363, 616. see also taiga bean goose;
model resolution, 106	tundra bean goose
ocean components, 106–107	beans, dry, 810, 811
"pole problem," 106–107	bear ( <i>Ursus</i> spp.), 301, 549, 609, 881, 953. see also individual
transient response experiments, 104	species
uncertainties in projections, 120	bearded seal (Erignathus barbatus), 488, 624
and vulnerability assessments, 958–959	in Bering Sea, 752
atmospheric circulation, 23–26	harvest by indigenous peoples, 652, 667
and ozone levels, 154, 157	harvest of, 622, 625, 627, 628, 632
and pollutant transport, 954	projected impacts, 504, 510, 1011
atmospheric general circulation model (AGCM), 102–103	Beaufort Gyre, 44, 458, 467, 471, 473, 474, 475
Atmospheric Model Intercomparison Project (AMIP), 108	Beaufort Sea, 476–477, 493, 506
atmospheric pressure, 24–26	beaver (Castor canadensis), 363, 654, 879, 880, 881, 897
attribution, 42	beech (Fagus sylvatica), 568, 571
observational data for model evaluation, 116, 117	beetles, 256, 549, 552, 553, 1016
observed trends, 23, 42	Beijerinckia indica, 260
observed trends compared to models, 42	beluga whale (Delphinapterus leucas), 489
projected change, 127, 128, 134–135	in Bering Sea, 752
attribution of cause of climate change, 3, 105, 909, 991	harvest by indigenous peoples, 652, 654, 669, 753
aufeis, 208, 358	harvest management in Canada, 612, 627
auks, 504, 626, 627, 653. see also great auk; little auk	harvest of, 622–623, 625, 629, 633
Aulacomnium turgidum, 258	historic change, 663
autumnal moth (Epirrita autumnata), 268, 280, 281, 299, 300,	importance of sea ice, 659
549–550, 557–558, 834	indigenous observations of change, 70
avalanches, 200, 869, 917, 918–919	migration, 546
avian influenza, 881, 882	overharvesting, 487
Aythya affinis. see lesser scaup	projected impacts, 380, 504, 510–511, 759, 1013
Aythya collaris. see ring-necked duck	tracking, 631
Aythya marila. see greater scaup (Aythya marila)	Bering land bridge, 252
Aythya valisineria. see canvasback duck	Bering Sea. see also marine Arctic
Azotobacter, 260	climate regime shifts, 753–754, 1018
n	commercial fisheries, 746–770
B	"Donut hole," 747, 750, 765
Bacillus, 260, 283	ecosystem description, 747
Back Basin, 227 bacteria, 260, 303, 312–313, 479, 481–482, 540, 544. see also	factors affecting ecosystem, 544 high-seas fishing moratorium, 694
microorganisms	nutrient levels, 491
Baffin Bay, 476–477, <i>493</i>	observed ecosystem change, 659, 1009–1010
Baker Lake	"oscillating control hypothesis," 755
indigenous knowledge case study, 82–84	past climate variability and fish, 501–504
indigenous observations of change, 69, 71	productivity, 493, 754–755
water level, 83–84	projected impacts, 476–477, 757–761, 1010
Balaena mysticetus. see bowhead whale	variability of temperature and salinity, 469
Balaenoptera acutorostrata. see minke whale	Bering Shelf, 480–481, 483, 491–493, 506–507
Balaenoptera borealis. see sei whale	Bering Strait, 28, 69, 71
Balaenoptera musculus. see blue whale	Beringia, 259, 328, 332, 584, 997, 998, 1016. see also Region 3
Balaenoptera physalus. see fin whale	berries, 796, 797
Balanus balanoides, 483	Berryteuthis magister magister. see commander squid
Balanus balanus, 483	Betula exilis. see dwarf birch
balsam poplar (Populus balsamifera), 252, 836	Betula nana. see dwarf birch
Baltin, 653	Betula papyrifera. see paper birch
bar-tailed godwit (Limosa lapponica), 258	Betula pendula. see silver birch
Barents Sea	Betula pubescens ssp. czerepanovii. see mountain birch

Betula spp. see birch	observed change in distribution, 287
Beverly-Qamanirjuaq Caribou Management Board, 612-613, 637,	overharvesting, 601
960	projected change in distribution, 287, 290–291
bilberry. see blueberry	projected impacts, 660
biodiversity, 11, 543, 1015–1018	and UV radiation, 268
of animals, 259–260	warming and disease, 880 birds, aquatic. see also birds, migratory; waders
definition, 541 in Finland, 795, 1016	and avian influenza, 881, 882
of freshwater ecosystems, 358, 998	breeding and resource availability, 418
genetic diversity and adaptation, 583	and contaminant transport, 431
and genetic variability, 582	diversity, 259
human impacts on, 560–567	harvest by indigenous peoples, 654
of human-modified areas, 554–555	harvest in Russia, 609
in Icelandic forests, 799	and introduced predators, 605
impacts of change in UV radiation, 424, 425	observed habitat change, 897
impacts of land and water management, 562–564	potential decoupling of environmental cues, 437
impacts on biogeochemistry, 310–311	predation pressure and climate change, 301
importance to indigenous peoples, 540 indicators, 585, 588–589	projected habitat loss, 288–289
influence of seasonal change, 543	projected impacts, 417–419, 436, 604, 659, 660 protected areas, 602
of lakes, 362–363	birds, migratory, 247, 252
latitudinal gradients, 256–260, 605, 606, 1016	and avian influenza, 882
of lichens, 273	and boreal forests, 785–786
of microorganisms, 260–262	breeding areas, 1005, 1006
monitoring change in, 579, 585–589	and disease transmission, 881
paleoecological change, 584	diversity, 259
of plants, 255–259	impacts of human activities, 564
pollution impacts, 564–566	management and conservation needs, 599
projected impacts, 263, 326, 328–329, 384–386, 418, 435–436,	and multiple stresses, 290–291
567–579, 588–589, 997, 998, 1012	observed change in arrival timing, 571
projected impacts on genetic diversity, 572–574 in Region 1, 1005	projected impacts, 267, 574–575, 1017 threats to, 581
research and monitoring recommendations, 580–589	and West Nile virus, 880
of rivers, 359	and wetland drainage, 562
sensitivity to climate change, 247	wildlife management challenges, 601–602
of soils, 540	bison (Bison bison), 80, 881
of Svalbard, 1016	bivalves, 483
of terrestrial ecosystems, 246, 255–263	black bear (Ursus americanus), 619
of tundra and polar desert ecosystems, 548-549	black-bellied plover (Pluvialis squatarola), 258, 361
of wetlands, 364	black brent. see Brent goose (Branta bernicla)
biodiversity conservation	black cottonwood (Populus trichocarpa), 800
approaches, 541, 558, 559–560, 579–591, 602–605	black-footed albatross ( <i>Phoebastria nigripes</i> ), 635 black guillemot ( <i>Cepphus grylle</i> ), 480, 489, 627, 629–630, 635
freshwater ecosystems, 547 genetic diversity, 572–574	black-headed gull ( <i>Larus ridibundus</i> ), 626, 629–630, 630
and indigenous knowledge, 558–559	black-legged kittiwake ( <i>Rissa tridactyla</i> ), 489, 629–631, 633, 635,
marine key issues, 546	1009–1010
modeling change, 568, 569	black scoter (Melanitta nigra), 86, 630
projected impacts, 541	black spruce (Picea mariana), 370
and protected areas, 605	in Alaska, 826–828
research and monitoring recommendations, 580-589	albedo, 789
species conservation, 556–558	in Canada, 792, 826–828, 850–851
strategic issues, 541–542	carbon dioxide fertilization, 845
biological diversity. see biodiversity	carbon storage, 843
birch (Betula spp.), 262, 293, 551, 784. see also individual species	observed change, 832
defoliation by autumnal moth, 299, 300 impacts of UV radiation, 852	projected impacts, 828 regeneration and temperature, 851
indigenous observations of change, 685	regeneration and wildfire, 836
past distribution, 368–369	temperature response, 826–828
projected change in distribution, 584	and wildfires, 838, 840
regeneration, 836	black-tailed godwit (Limosa limosa), 287
birds. see also seabirds; waders; wildlife; individual species	blackfishes (Dallia spp.), 397
cavity-nesting insectivorous, 797	blue-green algae. see cyanobacteria
change in nesting timing, 571	blue mussel (Mytilus edulis), 483, 504, 716, 739
and contaminants, 600	blue tit (Parus caeruleus), 1006
diversity, 259–260, 1016	blue whale (Balaenoptera musculus), 489
diversity and latitude, 258	blue whiting (Micromesistius poutassou), 508, 700, 714–715
diversity and temperature, 256	blueberry (Vaccinium myrtillus), 273, 275, 276, 276, 277, 278,
endangered status, 558 harvest of, 247	568, 571, 654, 792 bluefin tuna ( <i>Thunnus thynnus</i> ), 500, 508, 732, 740
importance to indigenous peoples, 599	bluegill (Lepomis macrochirus), 398
indigenous observations of change, 70–71, 86, 89, 668, 993	bluntnose sixgill shark (Notidanus griseus), 500
number of species 556 580	has hilherry see has whartleherry

bog blueberry. see bog whortleberry	Brosme brosme. see tusk
bog whortleberry (Vaccinium uliginosum), 271, 276, 277, 571	brown bear (Ursus arctos), 252, 258, 545, 1016
Bombus spp. see bumblebees	in Fennoscandia, 616, 977
Bonn Convention. see Convention on the Conservation of	harvest by indigenous peoples, 653
Migratory Species of Wild Animals	indigenous observations of change, 70
Boreal Ecosystem-Atmosphere Study (BOREAS), 789, 826, 828,	in Norway, 616
845	population monitoring, 620
boreal forest ecosystems	population status in Alaska, 619
albedo, 785	brown lemming (Lemmus sibiricus), 258
arctic importance, 786–787	brown trout (Salmo trutta), 382, 403, 407, 409, 410, 412, 563,
and beaver, 879	1012–1013
biodiversity, 790	brucellosis, 880, 881, 1011
biodiversity conservation, 551–554, 789	Brünnich's guillemot. see thick-billed murre
and carbon dioxide fertilization, 845–846	bryophytes, 552, 580, 1016
carbon storage, 305, 785, 791, 841, 844–845	bryozoans, 483
characteristics, 292	Bryum cyclophyllum, 258
climate projections, 802–806	buff-breasted sandpiper ( <i>Tryngites subruficollis</i> ), 557
climate projections, approach, 801–802	buffalo. see bison
climate variability, 788–789, 803–804	buildings. see infrastructure
climatic features, 787–788	bull trout (Salvelinus confluentus), 402
defined, 784	bumblebees, 258
disease, 805	burbot (Lota lota), 378, 397, 412, 563, 654
distribution, 787	Buteo lagopus. see rough-legged buzzard
disturbance and carbon cycle, 841–842	buttercups ( <i>Ranunculus</i> spp.), 272. see also individual species
global importance, 784–786	butterflies, 256, 287, 571
growth response and climate, 820–830	buzzards, 71, 304. see also rough-legged buzzard
impacts of pollution, 565, 566	
impacts of UV radiation, 851–853	C
influence on climate, 789	cadmium, 952, 954, 964, 979, 982. see also heavy metals
insect disturbances, 576, 792, 794, 804, 805, 830–835	Cajander larch (Larix cajanderi), 815–817
land tenure and management, 789–801	Calamagrostis lapponica, 295
latitudinal gradients of growth response, 821–824	calanoid copepods, 711–712, 732, 754–755
management and adaptation, 805–806	Calanus spp., 494
management and carbon balance, 785	impacts of past climate variability, 497, 498
multiple stresses on, 797	impacts of UV radiation, 513, 515, 516
and nitrogen availability, 845–846	as indicators of climate change, 482
observed change, 1009	as prey, 507, 511, 512, 721
projected change in distribution, 321, 326, 327–328, 569–570,	projected impacts, 504, 506
999, 1005–1006, 1007–1008, 1012	Calcarius lapponicus. see Lapland longspur
projected impacts, 551–554, 570, 789, 792–794, 797, 802–806,	Calidris acuminata. see sharp-tailed sandpiper
821, 822–824, 831, 845–846, 1009	Calidris alba. see sanderling
rate of migration, 321	Calidris alpina. see dunlin
research recommendations, 854	Calidris canutus. see red knot
role in freshwater discharge, 786	Calidris ferruginea. see curlew sandpiper
surprise responses, 804–806	Calidris fuscicollis. see white-rumped sandpiper
and wildfires, 553, 835–841	Calidris maritima. see purple sandpiper
boreal woodlands. see tundra-taiga ecotone	Calidris mauri. see western sandpiper
Bosmina longirostris, 422	Calidris melanotos. see pectoral sandpiper
bottlenose dolphin (Tursiops truncatus), 629	Calidris minuta. see little stint
botulism, 867, 880	Calidris ruficollis. see red-necked stint
boundary layer, atmospheric, 106	Calidris rayicoms, see rea-necked still
bowhead whale (Balaena mysticetus), 489, 623–624	Calliopidae sp., 483
in Bering Sea, 752	Callorhinus ursinus. see northern fur seal
harvest by indigenous peoples, 652, 663, 672, 674, 753, 956	Calluna vulgaris, 275
harvest of, 627, 633	Canada. see also individual provinces
historic harvesting, 699, 701, 753	forest area, 784
impacts of past climate variability, 504	forest tenure and management, 791–793
overharvesting, 487, 560, 1016	observed change in glaciers and ice sheets, 204–205
projected impacts, 380, 504, 510-511, 1013	projected impacts on boreal forest, 792–793
Branta bernicla. see Brent goose	protected areas, 604
Branta bernicla bernicla. see dark-bellied brent goose	sanitation systems, 884
Branta bernicla hrota. see light-bellied brent goose	wildfire regimes and management, 837–838
Branta canadensis. see Canada goose	wildlife management, 610–615, 626–628
Branta leucopsis. see barnacle goose	Canada goose (Branta canadensis), 70-71, 288, 363, 574, 662
Branta ruficollis. see red-breasted goose	Canadian Archipelago, 476–477
Brassicaceae, 293	Canadian Climate Impacts and Adaptation Research Network
Brent goose (Branta bernicla), 288, 363, 630	(CCIARN), 79
brittle stars, 487, <i>561</i>	cancer, 880
broad whitefish (Coregonus nasus), 395, 399, 411, 436	Cancer magister. see Dungeness crab
bromine, 154, 155, 170	Canis lupus. see wolf
bronze birch borer (Agrilus anxius), 576, 834	canola, 807, 810, 852
brook trout (Salvelinus fontinalis), 398, 407, 409, 410, 1012	canvasback duck (Aythya valisineria), 363
, , , , , , , , , , , , , , , , , , ,	1 2 2 111111111111111111111111111111111

capacity building	recent change in terrestrial ecosystems, 316–317
and co-management of wildlife, 613	soil processes affecting, 313
and human health, 893	from terrestrial ecosystems, 308–309
for indigenous peoples, 686–687	from wildfires, 842
capelin (Mallotus villosus), 484	and winter respiration by microorganisms, 269–270
in Bering Sea, 747–748	carbon monoxide, 312, 313
in Central North Atlantic, 711–712	carbon storage, 1014
commercial fisheries, 697, 704, 707, 714, 734, 737–738, 748,	in boreal forest, 550, 785, 791, 830, 841
761	in Icelandic forests, 800
distribution, 484	and land use change, 844–845
food requirement, 484	in Norwegian forests, 798
harvest by indigenous peoples, 652	in peatlands, 841
impacts of past climate variability, 500, 501, 503, 717–719,	projected change, 323, 851, 1015
737–738, 739, 755, 766	and soil temperature, 820–821
impacts of temperature, 495, 496	in tundra, 34, 550
migrations, 711	and vegetation type, 842–844
in northeast Atlantic, 696	and wildfires, 835
in northwest Atlantic, 732	Cardamine pratensis, 258
overharvesting, 624	cardiovascular disease, 866, 869, 872, 880, 883, 894–895, 897,
past distribution change, 500	1000
as prey, 486, 488, 489, 504, 512, 696	Carduelis chloris. see greenfinch
productivity, 479, 484	Carduelis hornemanni. see Arctic redpoll
projected change in distribution, 545	Carex aquatilis, 285
projected impacts, 504, 507–508, 720–721, 740–741, 758, 759,	Carex bigelowii, 273, 285, 294, 582
768, 1007, 1013	Carex bigelowii/arctisiberica, 257, 258
capercaillie. see wood grouse	Carex chordorrhiza, 258
Carabus truncaticollis, 258	Carex duriuscula, 258
carbon balance	Carex auriuscuia, 238 Carex ensifolia, 582
approaches, 315–316	
	Carex lugens, 582
and arctic biodiversity, 541	Carex spp., 285, 311, 317, 386 Carex stans, 257, 258, 285, 294, 582
Arctic Ocean storage, 34	
of arctic wetlands, 364–365, 434–435 and boreal forests, 785–786	Carex subspathacea, 300 caribou (Rangifer tarandus), see also International Porcupine
	, ,
carbon dioxide contribution, 317–318	Caribou Board; reindeer/caribou
and ecosystems, 34 and growing season timing, 274	barren-ground, 969 Bathurst herd, 611
impacts of recent climate change, 315–318 marine Arctic, 519–520	Beverly herd, 611, 620 Bluenose herd, 611
marine Arctic uncertainties, 521	and brucellosis, 880, 881
· · · · · · · · · · · · · · · · · · ·	early Holocene range, 51
marine ecosystems, 479	· · · · · · · · · · · · · · · · · · ·
methane contribution, 317–318	Grant's caribou, 969
and nutrient cycles, 846	harvest and late freeze-up, 75
and peatlands, 32	harvest by indigenous peoples, 654, 667, 669, 672, 674, 675,
and permafrost degradation, 319, 391–393, 434–435	680–685
projected impacts, 309–310, 316–318, 321–324, 325, 327–328,	harvest management in Canada, 612 historic range, 680–681
547, 998, 1004, 1006, 1012	C /
role of lichens, 563	impacts of freeze-thaw events, 660
and sea ice extent, 505	impacts of ice layers, 675
storage in terrestrial ecosystems, 305	impacts of late freeze-up, 75
and UV radiation, 310	impacts of oil and gas development, 621
and vegetation change, 999	impacts of projected change in sea ice, 659
and wildfires, 841–842, 843	importance to tourism, 600
carbon cycle	indigenous observations of change, 70–71, 73, 81, 280, 585,
alkalinity pump, 518	611, 615, 651, 660, 661, 677, 684–685, 993
biological pump, 518	management arrangements, 637, 644–648
and boreal forests, 841–846	management arrangements and indigenous subsistence
oceanic role, 515–519	activities, 664–665
physical pump, 515–518	migration cues, 67
and terrestrial ecosystems, 305–311	multiple stresses on, 610
carbon dioxide (CO <sub>2</sub> )	observed change in distribution, 287, 681
and boreal forests, 785	observed change on Banks Island, 666–667
contribution to carbon balance, 317–318	paleoecology, 250
fertilization effect, 568, 813, 845–846	Peary caribou, 570, 613, 614–615, 675, 969, 1012
and ozone chemistry, 155	and permafrost degradation, 883
and permafrost degradation, 459	population monitoring, 620
plant response, 265, 275–276, 299	population status in Alaska, 619
soil sources and sinks, 312	populations in Canada, 611
and stratospheric temperature, 155	Porcupine herd, 618, 620, 654, 681–685
carbon dioxide emissions	Porcupine herd management, 644–648
impacts of permafrost degradation, 391-393	projected impacts, 615, 645, 660, 684, 883, 1010, 1011, 1012
impacts of projected change in coastal permafrost, 219	Qamanirjuaq herd, 611, 620
impacts of projected change in lakes, 393	and spread of Giardia, 880

subspecies distribution, 681	in Alaska and Canada, 818-819
threats from hunting, 613–614	impacts on fish stocks, 699-700, 716-719, 736-737, 753-757
Western Arctic herd, 611, 618–620	climate, present-day, 10, 22-32, 244-245, 356-357
woodland, 969	climate data
carp bream (Abramis brama), 406–407	observational for model evaluation, 110–112
carrots, 808	reanalyses, 110–112
Caryophyllaceae, 293	climate models. see also atmosphere-ocean general circulation
Caspian seal, 625	model; general circulation model; regional climate
Cassin's auklet, 635	models
Cassiope spp., 262, 293	criteria for ACIA selection, 108-109
Cassiope tetragona, 247, 257, 258, 272, 273, 289, 302	described, 102–109
Castor canadensis. see beaver	equilibrium response experiments, 104
cataract, 877, 878, 899, 953, 1001	hierarchy of, 102–103
caterpillars, 268, 282	intercomparison projects, 108, 108–109
Catharacta skua. see great skua	research recommendations, 1019
cephalopods, 488, 489, 511, 758. see also squid	selection for ACIA, 108–109
Cepphus columba. see pigeon guillemot	uncertainties related to, 331–332
Cepphus grylle. see black guillemot	climate models, ACIA-designated, 8, 103
Cerastium beeringianum, 247	atmospheric pressure projections, 127, 128
Cerastium regelii, 258	and CMIP, 128–129
cerebral vascular disease, 872	evapotranspiration projections, 187–188
Ceriodaphnia quadrangula, 422	features of, 110
Cervus elaphus. see red deer	geocryological hazard projections, 927–928
Cervus nippon. see Sika deer	ocean projections, 476
Cetorhinus maximus. see basking shark	permafrost zone projections, 211–215
Cetraria islandica, 258, 276	precipitation projections, 126–127, 128–129, 187–188, 918, 994
Cetrariella delisei, 258 Chaetozone setosa, 483	
char (Salvelinus spp.), 71, 83, 395, 397, 1010. see also individual	projections, use in assessment, 118–119, <i>122</i> scenarios, use for agriculture projections, 809–810
species	scenarios, use for agriculture projections, 809–810 scenarios, use for forest projections, 800–806
Charadrius semipalmatus. see semipalmated plover	scenarios of future climate, 119–129
chemical transport models, 106	scenarios of future commate, 119–129 scenarios of future vegetation, 570
Chernobyl, 565, 979	sea ice projections, 192–194, 195, 471, 472, 924–926
Chiloxanthus pilosus, 258	simulation of observed air temperature, 114, 115
chinook salmon (Oncorhynchus tshawytscha), 654, 752, 760	simulation of observed all temperature, 114, 113
Chionoecetes bairdi. see Tanner crab	simulation of observed cloudiness, 116–118
Chionoecetes opilio. see snow crab	simulation of observed erotatiness, 110–116
Chipewyan, 680	simulation of observed sea-ice extent, 118
chironomids, 364, 365–366, 564	simulation of observed sea-level pressure, 116, 117
Chlamys islandica. see Iceland scallop	simulation of observed snow cover, 118
chlorinated dioxins and furans, 565	simulation of observed solar radiation, 117, 118
chlorine, 154–156, 170	snow cover projections, 199–200, 201, 918
chlorofluorocarbons (CFCs), 154, 155, 994	temperature projections, 121–125, 128–129, 913–914, 918,
chlorophyll-a, 515, 520	994
Chone paucibranchiata, 483	climate observations
Choristoneura fumiferana. see eastern spruce budworm; spruce	history, 34–35
budworm	limitations of, 22, 31, 35, 39
Chrosomus eos. see northern redbelly dace	station density, 34–35
Chrysaora melamaster, 497	Climate of the 20th Century project, 108–109
chrysophytes, 481	climate scenarios
Chrysosplenium alternifolium, 258	ACIA climate baseline, 8, 112–113
Chukchi peoples, 626, 652, 968	ACIA model projections, 119–129, 373–376
Chukchi Sea, 381, 476–477, 481, 483, 491, 492–493	ACIA time slices, 8
Chukotka, Russia, 609, 610, 625, 1016	analogue, 8, 101
chum salmon (Oncorhynchus keta), 71, 75, 752, 760	defined, 7, 101
ciliates, 480, 481–482	improving, 140–144
Cinclidium arcticum, 258	selection of, 7–8, 102–103
Circumpolar Active Layer Monitoring Program (CALM), 216,	synthetic, 101
319, 676, 910	types of, 7–8
Circumpolar Biodiversity Monitoring Program, 585, 588–589	uncertainties in model projections, 120
Circumpolar Protected Area Network (CPAN), 541, 559, 587,	climate variability
589–590, 603	and biodiversity, 384–385
cisco (Coregonus spp.), 654. see also whitefish; individual species	in boreal forest region, 788–789
Cladina spp., 258, 270	and caribou migration routes, 610
Cladocera, 365–366	in climate model projections, 803–804
Cladonia spp., 247, 270, 276	and glacier mass balance, 206
clams, 74–75, 483, 487, <i>504</i>	impacts on Arctic char, 400
Clangula hyemalis. see long-tailed duck	impacts on Atlantic cod, 718
Clethrionomys rufocanus. see grey-sided vole	impacts on fish, 412, 414, 494, 497, 500, 503, 507
Clethrionomys spp., 549	impacts on fisheries, 412, 416, 417, 767
Climacium dendroides, 258	impacts on indigenous peoples, 665, 668, 669, 676, 685, 963,
climate, past, 788–789. see also paleoclimate	1001

impacts on marine ecosystems, 659, 740	historic change, 663–664
impacts on reindeer, 967	impacts of past climate variability, 500
impacts on reindeer husbandry, 970, 974-976	impacts of UV radiation, 516
impacts on ungulates, 676	marine biomass, 479
indigenous observations of change, 74, 81, 82–83, 85–90, 88,	marine productivity, 479
94, 660, 685	projected impacts, 504, 508, 770–771, 1010
and model projections, 100, 103–105, 109, 112, 119, 120, 124,	Coelodonta antiquitatis. see woolly rhinoceros
134, 140, 143–144	coho salmon (Oncorhynchus kisutch), 654, 752, 760
modes of, 24–26	cold halocline, 30, 45
and ozone depletion, 156–157, 995	cold urticaria, 873
paleoclimatic, 46–54, 819	cold waves, 119
and reindeer husbandry, 658	collared lemming (Dicrostonyx groenlandicus), 249, 251,
20th century, 34–46	266–267, 301
climate warming. see temperature, air	collembolans, 260, 268, 296, 540, 580, 1016
Clostridium, 260	diversity of, 549, 1016
cloudberry (Rubus chamaemorus), 88, 271–272, 288, 654	colored dissolved organic matter, 370, 373, 419–425, 420, 421
clouds	Commander Islands, 605, 626
climate model representation, 106	commander squid (Berryteuthis magister magister), 761
and DMS, 143	commercial activities
feedback, 994	projected impacts, 200
impacts of projected change in sea ice, 195	common dace (Leuciscus leuciscus baicalensis), 406
impacts on plants, 277–278	common eider (Somateria mollissima), 363, 489, 573, 617, 626,
and marine ecosystems, 520	627, 630, 633, 635, 653
observational data for model evaluation, 116–118	common frog (Rana temporaria), 422
projected change, 127, 470	common guillemot. see common murre
and UV radiation, 153, 162, 166–167, 174, 278, 995, 996	common loon (Gavia immer), 630, 633, 635
clover, 807, 810	common merganser (Mergus merganser), 629
Clovis hunters, 252–253	common murre ( <i>Uria aalge</i> ), 489, 572, 623, 629, 630, 633, 635,
Clupea harengus. see Atlantic herring	1009
Clupea pallasi. see Pacific herring	common snipe (Gallinago gallinago), 287
Clyde River, Nunavut, Canada, 664	common teal (Anas crecca), 258
impacts case study, 671–675	common tern (Sterna hirundo), 630
indigenous knowledge case study, 82–84	Conservation of Arctic Flora and Fauna (CAFF), 6, 541, 559–560
indigenous observations of change, 69, 70	600, 623. see also Circumpolar Biodiversity Monitoring
co-management. see wildlife management	Program; Circumpolar Protected Area Network (CPAN
coal fish. see saithe	contaminants. see also heavy metals; persistent organic pollutants
coal mining. see mineral extraction	specific chemical species
coastal communities	in arctic food chains, 600, 640, 867, 964
and change in fish stocks, 561	and biodiversity, 1016
economic importance of fisheries, 763–764	human health impacts, 890–892, 953–954
and erosion, 921, 940, 1010–1011	impacts on biodiversity, 564–566
erosion protection, 923	impacts on boreal forests, 795, 797, 798, 1017–1018
impacts of observed change, 660	impacts on indigenous peoples, 678
multiple stresses on, 771	impacts on reindeer, 657
observed change, 661–662	impacts on wildlife, 615, 620
and permafrost degradation, 897	and increased precipitation, 545
1 0	1 1
projected impacts, 195, 208, 219, 235–236, 659, 767, 1004,	in marine food chain, 624
1014	in marine wildlife, 602
vulnerability to sea-level rise, 231	monitoring needs, 640
coastal erosion, 659	projected impacts, 412, 415, 427–433, 436–437, 566, 947,
and carbon balance, 518–519	954–955, 1011
contributing factors, 233–234, 920–921	research recommendations, 982
and late freeze-up, 75	in traditional foods, 654, 657
at Nelson Lagoon, Alaska, 77	transport pathways, 890–892
observed change, 897, 921-922, 963, 999, 1010-1011	and UV radiation, 424
and permafrost degradation, 217, 219	Convention on Biological Diversity, 541, 558–559, 572, 582, 590
projected impacts, 235–236, 922–923, 927, 952, 999, 1004,	604, 956
1009, 1013	Convention on the Conservation of Migratory Species of Wild
responses to, 93	Animals, 564, 574, 630
and sea ice, 76–77, 232–233	copepods, 480, 481–483, 489, 493–494, 512, 513, 700, 758
and sea-level rise, 231	coping capacity, 981, 1014
at Shishmaref, Alaska, 888–889	constraints on, 976–980
St. George Island, 77	of fisheries, 708–709, 729–730, 745
in Tuktoyaktuk, Canada, 940	of indigenous peoples, 665
	· · · · · · · · · · · · · · · · · · ·
at Varandei oil storage facility, Russia, 940	of reindeer husbandry, 974–976
and wetlands, 998	coping mechanisms
coastal management, 559, 601	definition, 955
coastal stability. see coastal erosion	coping strategies, 1020
cod. see also Arctic cod; Atlantic cod; Greenland cod; Pacific cod;	and human health, 900
Pacific tomcod; polar cod	for indigenous peoples, 666, 955–957
aquaculture, 703, 708, 716, 765	in Sachs Harbour, NWT, 668–670, 963
food requirement, 484	and vulnerability analysis, 957

copper. see mineral extraction	Dansgaard-Oeschger events, 49–50
coral reefs, 561, 564	Daphnia spp., 387, 388, 422, 425, 514, 564
Corallorrhiza spp., 258	dark-bellied brent goose (Branta bernicla bernicla), 574
Coregonus. see whitefish	dark-leaved willow (Salix myrsinifolia), 853
Coregonus albula. see vendace	DDT. see persistent organic pollutants
Coregonus artedi. see lake cisco	deepwater prawn (Pandalopsis dispar), 483
Coregonus autumnalis. see Arctic cisco	deepwater shrimp. see northern shrimp
Coregonus clupeaformis. see lake whitefish (Coregonus	deforestation, 1017–1018
clupeaformis)	and carbon dioxide emissions, 844
Coregonus nasus. see broad whitefish	impacts on indigenous peoples, 678
Coregonus pidschian. see Siberian whitefish	impacts on reindeer pastures, 657
Coregonus sardinella. see least cisco	by reindeer overgrazing, 680
cormorant ( <i>Phalacrocorax</i> spp.), 627, 633. see also double-crested	and tundra, 247, 320, 1005
cormorant; great cormorant; Pallas's cormorant; pelagic	and wind speed, 324
cormorant; red-faced cormorant	Delphinapterus leucas. see beluga whale
corneal damage, 996	deltas, 1016. see also estuaries
Cortusa matthioli, 258	ecosystems of, 360–361
Corvus corax. see raven	impacts of sea-level rise, 235
Cotesia sp., 550	projected impacts, 360, 379–381, 418, 434
cottongrass ( <i>Eriophorum</i> spp.), 257, 258, 262, 273, 275, 278, 293,	Denali National Park, 567
311, 317, 549	Dendroctonus rufipennis. see spruce bark beetle
Cottus cognatus. see slimy sculpin	Dendrolimus sibiricus. see Siberian silkworm
Couesius plumbeus. see lake chub	Dene, 599, 652, 664, 680, 681
cougar (Puma concolor), 78	observations of change, 1014
Coupled Model Intercomparison Project (CMIP), 108, 119,	subsistence activities, 653
128–129, 193	Dene Nation
crabs. see also Dungeness crab; king crab; red king crab; snow	indigenous knowledge case study, 78–82
crab; Tanner crab	vulnerability, 962
in Bering Sea, 746	Denendeh, 78–82. see also Dene Nation
commercial fishery, 742, 762, 763, 764, 766	Denendeh Environmental Working Group, 78–82, 962
historic harvesting, 762	dental cavities, 883
impacts of past climate variability, 503, 756–757, 766	diabetes, 866, 880, 883, 895, 897, 1000
management regimes, 765	diamond mining. see mineral extraction
predation by Atlantic cod, 739	diatoms, 365–366, 481, 506
projected impacts, 504, 760–761, 768	Dicentrachus labrax. see sea bass
cranberry (Oxycoccux spp.), 654	Dicranoweisia crispula, 258
Cree, 662, 680, 681	
	Dicrostonyx groenlandicus. see collared lemming
creep, 909, 914, 916	Dicrostonyx spp. see lemmings
creeping alkaligrass (Puccinellia phryganodes), 300	Dicrostonyx torquatus. see Arctic lemming
creeping juniper (Juniperus communis), 799	dimethyl sulfide (DMS), 143, 312
crested auklet (Aethia cristatella), 633, 635	dinoflagellates, 481
crossbills ( <i>Loxia</i> spp.), 267	Dioryctria reniculelloides. see spruce coneworm
crowberry (Empetrum nigrum), 273	disease. see also human health
crows, 71	and emigration of new species, 386, 395, 419, 436
crustaceans, 483, 488, 504, 512, 513, 540, 580, 754. see also	in human populations, 866–869
barnacles; copepods; Daphnia; krill; shrimp	impacts on biodiversity, 588
cryosphere, 22, 184. see also glaciers and ice sheets; ice, river and	projected impacts, 413, 415, 417, 419, 436, 879–881, 999
lake; permafrost; sea ice; snow	1004
cryoturbation, 210, 215	and sanitation infrastructure, 886-888
Cryptococcus laurentii, 262	and traditional foods, 867
cryptogams, 256	and UV radiation, 878
cryptophytes, 481	zoonotic, 880–881
cryptosporidium, 880, 881, 884	dissolved inorganic carbon (DIC), 515, 518
cultural diversity. see arctic residents; indigenous peoples	dissolved organic carbon (DOC)
	` , ,
cultures. see arctic residents; indigenous peoples	and carbon cycle, 518
curlew sandpiper (Calidris ferruginea), 258, 288, 290	impacts of change in, 390–391, 393, 420, 434
Curtonotus alpinus, 258	projected change in, 377–379, 420, 423
cyanobacteria, 262, 271, 275, 284, 481. see also microorganisms	and UV radiation exposure, 515, 516, 520
Cyclops scutifer, 422	dissolved oxygen (DO), 379
Cygnus columbianus. see tundra swan	diurnal temperature range, 119
Cyperaceae, 293	Dogrib, 680
Cystophora cristata. see hooded seal	Dolgans, 599
	Dolly Varden (Salvelinus malma), 75
D	anadromous behavior, 409
Dactylina madreporiformis, 258	importance of estuaries, 361
Dactylina ramulosa, 258	importance of groundwater, 402
Dahurian larch, 292	projected impacts, 385, 410, 411
Dall sheep (Ovis canadensis dalli), 654	thermal preferences, 395
Dallia spp. see blackfishes	double-crested cormorant, 635
Dall's porpoise ( <i>Phocoenoides dalli</i> ), 489, 752, 759, 765	dovekie. see little auk
dams. see impoundment	and the second water
dandelion (Taraxacum officinale), 605	
annation ( I ai anacain officiane), 000	

downscaling	energy balance, 314
of model-projected arctic change, 137–140	global links, 247
and regional climate modeling, 137	human impacts on, 253
research recommendations, 982	impacts of experimental manipulation, 295–298
statistical methods, 103, 136–137	impacts of poaching, 679
for vulnerability assessments, 958–959	importance to humans, 247
downy birch (Betula pubescens), 557–558, 562, 563, 795, 799–800	latitudinal variation, 292–295
Draba spp., 258, 293	prehistoric human impacts, 252
Dreissena polymorpha. see zebra mussel	projected impacts, 253–254, 998
Drepanocladus intermedius, 258	surprise responses, 249
drought	trophic interactions, 298–303 water balance, 314
adaptations of microorganisms, 270 and agriculture, 812	eel (Anguilla anguilla), 410, 544
and arctic animals, 281, 296	eider, 617. see also common eider; king eider; spectacled eider;
and arctic plants, 272, 302	Steller's eider
and boreal ecosystems, 552, 783, 826, 829, 1001	harvest of, 626, 627, 629, 631, 633, 634–635
and eastern spruce budworm, 833	impacts of sea level rise, 636
and human health, 881, 885	indigenous observations of, 77
and migratory birds, 290	management agreements, 623
paleoclimatic, 466	overharvesting, 631
projected change, 869, 999	El Niño-Southern Oscillation (ENSO), 3
and sensible heat flux, 318	and caribou behavior, 287
and spruce bark beetle, 832–833	and fish recruitment, 503
and wildfires, 679, 835	impacts on aquaculture, 703
Dryas integrifolia. see Arctic dryad	impacts on walleye pollock, 756
Dryas octopetala/punctata. see mountain avens	and infectious diseases, 879–880
<i>Dryas</i> spp., 262, 293, 294–295, 549	and lake water temperature, 429
duck plague, 880	and sea ice variability, 469
ducks, 358, 436, 490, 616–617, 654, 659, 668. see also individual	and sea otter populations, 632–633
species	and wildfires, 836–837
duckweed (Lemna spp.), 89, 363	Eleginus gracilis. see saffron cod
Dungeness crab (Cancer magister), 483	Elim, Alaska, 68
dunlin (Calidris alpina), 251, 258, 288, 361	Emberiza pusilla. see little bunting
Dupontia spp., 247, 258, 311	emerald shiner (Notropis atherinoides), 398, 405
dwarf birch (Betula exilis), 257, 258 dwarf birch (Betula nana), 247, 257, 258, 273, 274, 295, 299, 799	Emiliana huxleyi, 481. see also phytoplankton
dwarf Siberian pine ( <i>Pinus pumila</i> ), 293	emissions scenario, 119–121, 958, 993–994 A2, described, 8
dwarf willow (Salix herbacea), 265	A2, temperature projections using, 119
dwarf willow (Ballx herbacea), 203	A2 and B2 compared, 806
E	ACIA rationale, 120
earthworms, 580	B2, described, 8
East Siberian Sea, 493, 506, 518	B2, temperature projections using, 119
eastern spruce budworm (Choristoneura fumiferana), 833–834	defined, 7
echinococcus, 880, 881, 1011	IPCC IS92 scenarios, 119–121
echiuran worms, 487	IPCC Special Report on Emissions Scenarios, 7-8, 109, 109,
economic impacts, 4, 1013. see also agriculture; arctic residents;	119–121
fisheries; forestry; indigenous peoples; mineral	selection of, 7–8
extraction; natural resources; oil and gas development;	emperor goose (Anser canagicus), 288, 289
recreation; resource development; tourism	Empetrum hermaphroditum. see mountain crowberry
of projected change in fisheries, 416–417	Empetrum nigrum. see crowberry
research recommendations, 1019	Empetrum spp., 293, 294
ecosystems. see also boreal forest ecosystems; freshwater	enchytraeid worms, 540, 580
ecosystems; marine ecosystems; polar desert	endangered species
ecosystems; terrestrial ecosystems; tundra ecosystems;	bowhead whale, 753
wildlife management	definitions, 581–582
conservation, 543–560, 602–605, 651	Eskimo curlew, 363
decomposition in, 540–541 fragmentation and protected areas, 577–578	monitoring, 587 Red Lists, 541, 585, 624
habitat groupings, 543–544	spectacled eider, 363
impacts of UV radiation, 995	Steller sea lion, 753
multiple stresses, 605	Steller's eider, 363
novel, 550, 569, 570, 571, 789, 822–823, 851, 998, 1017	and undisturbed forests, 790, 795, 797
photosynthesis in, 540–541	energy balance
processes within, 540–541	of boreal forest, 789
projected impacts, 4, 215, 606	of terrestrial ecosystems, 314, 318–319
ecosystems, terrestrial. see also boreal forest ecosystems;	and vegetation distribution, 324
freshwater ecosystems; polar desert ecosystems; tundra	energy budget, surface
ecosystems	impacts of projected change in snow cover, 200
approach for assessment, 248-249	importance of mosses, 273
biogeochemical cycling, 305–311	importance of sea-ice albedo, 190
carbon balance, 315–318	energy consumption
disturbances, 245, 247	projected change in heating requirements, 936, 937

Enets people, 679	in vulnerability analysis, 959
engineering. see infrastructure	and water supply, 885–886
Engraulis mordax. see northern anchovy	Exxon Valdez, 565
Enhydra lutris. see sea otter	Eyak people, 656
epiphytic organisms, 480	eye irritations, 63
Epirrita autumnata. see autumnal moth	
epishelf lake, 381	F
equilibrium response, 104	Fagus sylvatica. see beech
Equisetum spp. see horsetail	Falco peregrinus. see peregrine falcon
Eremophila alpestris. see shore lark	Falco rusticolus. see gyrfalcon
Erignathus barbatus. see bearded seal	falcon, 71. see also gyrfalcon; peregrine falcon
Eriophorum spp. see cottongrass	Faroe Islands, 868, 869
Eritrichium nanum, 249	fast ice. see land-fast ice
ermine (Mustela erminea), 247, 258, 259, 301, 607	feedbacks, 7, 23, 990, 994, 998, 999, 1015
erosion. see also coastal erosion	albedo-temperature, 3, 196, 252, 315
impacts of permafrost change, 659, 913	and ecosystem processes, 315, 321–324, 550
and infrastructure, 923	ecosystems and greenhouse gases, 11
and landslides, 917–918	freshwater flux and thermohaline circulation, 315
projected change, 917	and glacial cycles, 47
and sanitation infrastructure, 888	greenhouse gas emissions and vegetation change, 315
and water supply, 886	ice-albedo, 33, 190
Eschrichtius robustus. see grey whale	methane hydrates, 315, 519
Eskimo curlew (Numenius borealis), 363	and ocean carbon processes, 518
Esox lucius. see northern pike	permafrost thawing, 459, 520
estuaries, 544	and rapid temperature change, 50
ecosystems of, 360–361	salinity, 478
nutrient limitation in, 491	sea ice-albedo, 459, 474, 476–477, 1005
projected impacts, 235, 379–381, 385, 418, 547, 923	soil carbon, 785
timing of spring phytoplankton bloom, 492	thermohaline, 459
ethics of climate change, 4, 64	vegetation-albedo, 785, 844
Eubalaena glacialis. see right whale	wildfire and carbon emissions, 842
Eubalaena japonica. see North Pacific right whale	wildfire and vegetation distribution, 840
Eucalanus bungii, 482	Fennoscandia
eulachon (Thaleichthys pacificus), 503, 755	forest tenure and management, 794–801
Eumetopias jubatus. see Steller sea lion	projected climate change, 971–972
Euphausia pacifica, 482	reindeer husbandry, 965
euphausiids. see krill	and Scots pine growth, 829–830
Euphrasia frigida, 263	wildfire regimes and management, 839–840
Eurasian oystercatcher (Haematopus ostralegus), 630	wildlife management, 616–617, 628–631
Eurasian wigeon (Anas penelope), 258	ferns, 580
European aspen (Populus tremula), 799	fieldfare (Turdus pilaris), 258, 260
European eel. see eel	fin whale (Balaenoptera physalus), 489, 615, 629, 652, 715, 732,
European perch (Perca fluviatilis), 405, 406	736, 753, 965
European pine sawfly (Neodiprion sertifer), 835, 1007	Finland
European shag ( <i>Phalacrocorax aristotelis</i> ), 629–630, 630	agriculture, 808
European storm petrel (Hydrobates pelagicus), 630	demographics, 868–869
European whitefish (Coregonus lavaretus), 395, 397, 411, 412,	forest area, 784
563, 576	forest tenure and management, 795–796
Eurynorhynchus pygmaeus. see spoon-billed sandpiper	indigenous observations of change, 69, 71
evaporation	projected impacts, 584, 795
measurement limitations, 185	projected impacts, 384, 773 protected areas, 541
present-day, 245	Finnmark, Norway
projected change, 126–127, 133–134	vulnerability case study, 962, 965–980
evapotranspiration	fire. see wildfires
measurement limitations, 31	firn, 31, 784
projected change, 812	firs (Abies spp.), 551. see also Siberian fir
for projecting agriculture impacts, 810	First Nation Council of Yukon Indians, 645
and vegetation change, 324	fish. see also fisheries; freshwater ecosystems; marine ecosystems
Even peoples, 652	individual species
Evenk peoples, 652	indigenous observations of change, 70–71, 993
Evenkiya, Russia, 609	as source of Vitamin D, 877
exclusive economic zone (EEZ), 694	warming and disease, 880 fish, anadromous, 544, 547. <i>see also</i> fish, diadromous; salmonids;
Exobasidium, 302	
exotic species. see non-native species	individual species
extreme events, 1018. see also drought; floods; storms	definition of, 356
and human health, 869–870, 881	indigenous harvesting, 747
impacts on aquaculture, 708	mercury concentrations in, 433
impacts on coosystems, 999	and nutrient/energy transfer, 361
impacts on freshwater fisheries, 413, 415, 416	observed change, 1010 projected impacts, 360–361, 402–403, 437, 998
indicators for, 899 modeling, 143–144	projected impacts, 300–301, 402–403, 437, 998
projections of, 144	

fish, catadromous, 544	in Greenland, 709–731, 964–965
definition of, 356	Greenland halibut, 697, 713, 714, 715, 733–734, 737, 748–749
projected impacts, 410	haddock, 698
fish, demersal. see groundfish	humpy shrimp, 749
fish, diadromous	in Iceland, 709–731
definition of, 356	impacts of past climate variability, 756
projected impacts, 393–406, 408–410, 436	impacts on arctic residents, 1000
species, number of arctic, 396–397	impacts on biodiversity, 555, 561, 564
thermal preferences, 394–399	impacts on indigenous communities, 657
fish, freshwater	impacts on indigenous harvesting, 559
adaptation to current climate, 357	impacts on marine mammals, 633
biodiversity, 547, 1016	invertebrates, 748
inventory needs, 580	the "Loophole," 496
mercury concentrations in, 405, 433	management and indigenous subsistence activities, 665
migratory timing change, impacts on fisheries, 413–414	management regimes, 694–695, 704–706, 707–708, 729, 742,
projected impacts, 393–408, 402–403, 412–413, 436, 437, 547,	746–747, 763, 765–766, 766–767, 770–771
570, 998	markets, 704, 764–765
projected range shifts, 588	multiple stresses on, 706–707
species, number of arctic, 11, 396–397	Newfoundland and Labrador, 731–746
thermal preferences, 394–399	in North Pacific and Bering Sea, 746–770
threats to migratory species, 581	in Northeast Atlantic, 695–709
fish, marine, 484–487	
	northern flathead sole, 751
adaptation to arctic environment, 557	northern shrimp, 697, 713, 714, 715, 722–723, 734–735,
and benthos, 483	738–739, 749
distribution and migration, 496	Pacific cod, 750–751
diversity, 1017	Pacific herring, 748
and human health, 897	Pacific salmon, 748, 752
impacts of climate factors, 494–496	pandalid shrimp, 749
impacts of oil pollution, 565	polar cod, 697, 748
impacts of past climate variability, 499–504, 699–700,	productivity of, 4
716–719, 736–739, 753–757	projected impacts, 739–741, 757–761, 770–771, 1001, 1002,
impacts of UV radiation, 514	1004, 1005, 1006–1007, 1010, 1013, 1017
inventory needs, 580	red king crab, 749
number of species, 484	redfish, 698, 714, 715
and ocean currents, 710–711	in Region 1, 1006
and ocean temperature, 520	research recommendations, 771–772
projected change in distribution, 569, 588, 694, 761	rock sole, 751
projected change in migration, 694	role in mixed economies, 656
projected impacts, 504, 507–508, 544–545, 570, 719–721,	roundfish, 748
739–741, 744–745, 757–761	shrimp, 742
reproductive cycle and temperature, 495	snow crab, 735, 749
temperature impacts on distribution, 496	Tanner crab, 749
threats to migratory species, 581	uncertainties in projections of change, 693-694
fish farming. see aquaculture	walleye pollock, 748, 749–750
fisheries	yellowfin sole, 751, 756
impacts of changing sea ice on harvests, 195	fisheries, freshwater and anadromous
impacts of projected change on indigenous peoples, 663–664	impacts on biodiversity, 555
impacts of UV radiation, 953, 999	and indigenous peoples, 558–559
and individual transferable quotas, 665	projected impacts, 410–416, 436
projected impacts, 999	projected impacts on specific fishery sectors, 416–417
fisheries, commercial, 990	value of, 411, 416–417
Alaska plaice, 751	Fisheries Joint Management Committee, 670
arrowtooth flounder, 748	flagellates, green, 481
Atlantic cod, 698–699, 707, 712–713, 714, 715, 722–723, 733,	Flakaliden experiment, 820–821
736–737, 742–744	flatfish. see also individual species
Atlantic herring, 696–698, 707, 714, 734	in Central North Atlantic, 711
in Barents and Norwegian Seas, 695–709	commercial fisheries, 748, 751
in the Bering Sea, 1009	impacts of past climate variability, 503, 756, 766
blue whiting, 714–715	projected change in distribution, 545
and boreal forests, 794	projected impacts, 504, 507–508, 758, 759, 760–761, 768
capelin, 714, 734, 737–738, 748	flathead sole (Hippoglossoides elassodon), 487, 502, 503
cod, 500, 696–697	flaw leads, 457
coping capacity, 709, 729–730, 768	flies, 540
	floods
crab, 742	
economic and social impacts on, 706–709, 721–725, 725–729,	factors affecting, 917
744–745, 766–768	impacts of projected change in freshwater runoff, 229
economic and social importance, 16, 700–704, 741–744,	impacts of projected change in river and lake ice, 223
761–766	impacts of projected precipitation change, 188
employment, 703–704	impacts of sea-level rise, 235–236
exclusive economic zones, 694	impacts of temperature change, 23, 32
and fish distribution, 496	and infrastructure, 917
flatfish, 748, 751	projected impacts, 917, 923, 997, 998, 1008, 1009

and rain-on-snow events, 43	biodiversity, 547
vulnerability analysis, 957	biodiversity conservation, 546–548
flounder (Pleuronectes ferrugineus), 652	biodiversity monitoring, 586
Folsomia spp., 260, 281, 282, 549	carbon storage, 1014
food chains, 540	characteristics, 1017
food webs, 540	and chlorinated dioxins and furans, 565
forest fires. see wildfires	and contaminants, 427–433, 565
forest tent caterpillar, 833–834	described, 11–12
1 ,	*
forest tundra. see tundra-taiga ecotone	historic change in, 365–372
forestry	impacts of industrial development, 677
in Alaska, 793–794	impacts of non-native species, 576
and biodiversity, 552–554, 560–561	impacts of projected change in freshwater runoff, 229
in Canada, 791–793	impacts of projected change in glaciers and ice sheets, 208
and carbon storage, 785–786	impacts of projected change in river and lake ice, 223
economic importance, 786	impacts of recent climate change, 371–372
and energy production, 796, 797, 799	impacts of UV radiation, 419–427, 947, 996
in Finland, 795–796	and mercury, 432–433, 565
in Iceland, 800	and PCBs, 565
impacts on biodiversity, 562–564	and persistent organic pollutants, 427–432
impacts on habitat, 616	projected impacts, 32, 372–438, 384–386, 433–437, 547–548,
impacts on indigenous peoples, 679–680, 685	997–998, 998, 1012–1013
impacts on reindeer overgrazing, 601	projected impacts on carbon dynamics, 391–393
in Norway, 798–799	projected impacts on fish, 393–410
projected impacts, 791, 1001, 1002–1003, 1007, 1008, 1010	projected impacts on food webs, 386–388
and protected areas, 604	projected impacts on hydro-ecology, 373–384
role in mixed economies, 656	projected impacts on production, 388–391, 434–436
in Russia, 790–791	projecting impacts, 372–438
in Sweden, 796–797	protected areas, 577
	•
forests, 247, 563, 784. see also boreal forest ecosystems;	research recommendations, 437–438
deforestation; forestry; treeline	structure and function of, 358–365
fork-tailed storm petrel, 635	and toxaphene, 565
Formitopsis rosea, 582	uncertainty in impacts projection, 354–356
Fort Chipewyan, Alberta, 654	water budgets, 357–358
Fort Smith, NWT, 654	freshwater hydrology, 31–32
fourhorn sculpin (Myoxocephalus quadricornis), 385	freshwater shrimp (Gammarus lacustris), 382
fox, 880, 881, 953. see also Arctic fox; red fox	fritillary butterfly, 586
Foxe Basin. see marine Arctic	frogs ( <i>Rana</i> spp.), 282, 422
Franz Josef Land, 206	frost days, 119
Fratercula arctica. see Atlantic puffin	frost heave, 909, 914–916, 930
Fratercula cirrhata. see tufted puffin	frost jacking, 909
Fratercula corniculata. see horned puffin	frostbite, 871–872
freeze-thaw events	Fucus spp., 483
and ecosystem diversity, 293	Fulmaris glacialis. see northern fulmar
impacts on animals, 142, 269, 279, 296, 301	fungi, 260–261, 298, 303, 540, 552, 571, 580, 797. see also
impacts on arctic species, 557	microorganisms
impacts on aspen, 833	Ç
impacts on caribou, 660	G
impacts on soil microorganisms, 297	Gadus macrocephalus. see Pacific cod
	Gadus macrocephatus, see Facilic cod
impacts on tundra vole, 280	
indigenous observations of change, 660	Gadus ogac. see Greenland cod
projected increase in, 200, 282	Galium densiflorum, 258
freezing rain	Gallinago gallinago. see common snipe
impacts on caribou, 279, 615, 661, 670, 883	Gambell, Alaska, 69
impacts on Collembola, 281	Gammarus lacustris. see freshwater shrimp
impacts on contaminant transport, 430	Gammarus oceanicus. 483
impacts on seabirds, 512	Gammarus setosus, 483
impacts on snow geese, 280	Gammarus wilkitzkii, 480, 482
indigenous observations of, 77	Gasterosteus aculeatus. see threespine stickleback
•	•
projected change, 200, 668, 952	gastroenteritis, 884
freshwater discharge. see also Arctic Ocean; river discharge	gastropods, 483
background, 223–225	Gavia adamsii. see yellow-billed loon
impacts of projected change, 228–229	Gavia arctica. see Arctic loon
measurement limitations, 225	Gavia immer. see common loon
observed trends, 225–227, 228	Gavia pacifica. see Pacific loon
projected change, 227–228, 229	Gavia stellata. see red-throated loon
research recommendations, 229–230	geese, 358, 490. see also individual species
role of boreal forests, 786	
	adaptive capacity, 291
and thermohaline circulation, 477	arctic-breeding species, 259
freshwater ecosystems	biodiversity monitoring, 586
and acidification, 565	habitat requirements, 268
adaptability to rapid climate change, 433	harvest by indigenous peoples, 654
barriers to species movement, 568	indigenous observations of change, 89

migration routes, 574	Great Bear Lake, 381
overharvesting, 560	great black-backed gull (Larus marinus), 629-631
projected impacts, 436, 574–575	great cormorant (Phalacrocorax carbo), 629–630
and snow duration, 280	great knot (Calidris tenuirostris), 288
general circulation model (GCM), 102-103. see also climate	great northern diver. see common loon
models	Great Salinity Anomaly, 45, 468, 717
Gentiana nivalis. see snow gentian	great skua (Catharacta skua), 288, 630
geography, 10	Great Slave Lake, 381
marine Arctic, 26	greater scaup (Aythya marila), 363
geometrid moths, 268, 281. see also autumnal moth; winter moth	greater shearwater (Puffinus gravis), 732
Geum, 278	greenfinch (Carduelis chloris), 1006
giant deer (Megaloceros giganteus), 250	greenhouse gases. see carbon dioxide; methane; nitrous oxide;
Giardia, 879, 880, 881, 884, 897	trace gases
glacial cycles, 47–50, 543. see also climate, past	Greenland
glacier buttercup (Ranunculus glacialis), 278	commercial fisheries, 709–731, 866
glaciers and ice sheets. see also Greenland Ice Sheet	demographics, 868–869
area by region, 202	economic importance of fisheries, 722–723, 724–725
background, 201–203	fisheries development, 663–664
and contaminants, 954	forest tenure and management, 799-801
dynamics and biodiversity, 550-551	human health, 894–895
equilibrium line defined, 31, 203	impacts of past climate change, 866
extent at last glacial maximum, 249	impacts on fisheries, 727–729
impacts of projected change, 208	marine ecosystem description, 710–711
impacts of temperature change on, 3	Norse settlement, 250, 562, 584
mass balance, 203	sanitation systems, 884
melting and landslides, 918	settlement, 799–800
melting and thermohaline circulation, 999–1000	social change, 865–866
observed change, 53, 997, 1009	vulnerability case study, 962–963, 964–965
observed change by region, 204–207	Greenland cod (Gadus ogac), 484, 486, 500
and precipitation, 31	Greenland halibut (Reinhardtius hippoglossoides), 486–487
projected impacts, 3, 119, 207–208, 659, 997, 1010	in Bering Sea, 748–749
research recommendations, 208–209	in Central North Atlantic, 711–712
and sea level rise, 994, 997	commercial fishery, 697, 713, 714, 715, 737, 748–749, 964
sensitivity to temperature change, 203	distribution, 484
and temperature, 31	harvest by indigenous peoples, 652
volume and sea-level equivalent, 201	historic fisheries, 733–734
as water source, 547	impacts of past climate variability, 503, 737, 756
glaucous gull ( <i>Larus hyperboreus</i> ), 361, 489, 566, 624, 629–630,	in northeast Atlantic, 696
635	in northwest Atlantic, 732
glaucous willow (Salix glauca), 278	past distribution change, 500
glaucous-winged gull (Larus glaucescens), 626, 635	as prey, 488
global climate, links to, 4–5, 12, 32–34, 55, 459, 541	projected change in distribution, 545
marine Arctic, 473, 519–520	projected enauge in distribution, 545 projected impacts, 504, 507–508, 740–741, 758, 759, 965,
sea ice, 23, 30	1007, 1013
thermohaline circulation, 32–33, 54	temperature and distribution, 496
global economy. <i>see also</i> economic impacts; fisheries; forestry;	Greenland Home Rule, 665, 724, 956, 964
indigenous peoples; mineral extraction; oil and gas	Greenland Ice Sheet, 201–202
development	area, 202
arctic contribution, 5, 16, 934, 937, 990, 1002, 1008	· · · · · · · · · · · · · · · · · · ·
and emissions scenarios, 8, 119–120, 332, 958, 993–994	freshwater storage, 31 impacts of projected change, 208
impacts on Arctic communities, 650–651, 771	
1	observed change, 205–206, 1012
impacts on Arctic fisheries, 708	prehistoric size, 46, 48
impacts on Arctic residents, 948, 950	projected change, 207, 997, 1004, 1012, 1018
impacts on indigenous peoples, 91, 657–658, 664–665, 685,	research recommendations, 208–209
955, 961, 1001, 1014	and sea-level rise, 203, 231
impacts on the Arctic, 636	sensitivity to temperature change, 203
Global Terrestrial Network for Permafrost (GTN-P), 209, 211,	simulation of observed change, 131–132
216, 910	Greenland right whale. see bowhead whale
Global Terrestrial Observing System (GTOS), 587	Greenland Sea, 493
global warming. see temperature, air	Greenland shark (Somniosus macrocephalus), 486
global warming potential (GWP), 318	Greenland turbot. see Greenland halibut
Globicephala melaena. see pilot whale	Greenland white-fronted goose (Anser albifrons flavirostris), 574
Glyptocephalus cynoglossus. see witch	grey phalarope. see red phalarope
Gmelin larch ( <i>Larix gmelinii</i> ), 815–816, 822–824	grey seal (Halichoerus grypus), 488, 504, 510, 622, 626–627, 628
gold. see mineral extraction	699, 715
goldeye (Hiodon alosoides), 396	grey-sided vole (Clethrionomys rufocanus), 280, 301
government	grey whale (Eschrichtius robustus), 489, 504, 510, 511, 624, 633,
role in addressing climate change, 78, 79, 91, 92, 640, 665,	652, 753
686–687, 900 grains, garael, 800, 812	greylag goose (Anser anser), 574, 617, 630
grains, cereal, 809, 812	grizzly bear. see brown bear
grazing, 562–563. see also Iceland; reindeer; reindeer husbandry	ground squirrel (Spermophilus parryii), 267, 653, 654
great auk (Pinguinus impennis), 12, 487, 560, 601, 1016	

groundfish	Haematopus ostralegus. see Eurasian oystercatcher
in Bering Sea, 746	hairy lousewort (Pedicularis hirsuta), 258
collapse of Newfoundland/Labrador fishery, 741–744	hairy willow (Salix lanata), 799
commercial fisheries, 710, 747, 762–764	halibut, 663, 708, 712, 765. see also Atlantic halibut; Greenland
fisheries and Steller sea lions, 747	halibut; Pacific halibut
historic harvesting, 712–713, 761–762	Halichoerus grypus. see grey seal
impacts of past climate variability, 503, 733, 739, 756	Haliotis rufuscens. see abalone
management regimes, 665, 699	halocline, 460
management under changing climate, 720, 766–767	halogens, 154–156, 170, 312. see also bromine; chlorine
monitoring, 502	harbour porpoise (Phocoena phocoena), 489, 629
past changes in abundance, 737	harbour seal ( <i>Phoca vitulina</i> ), 488
predators, 741	in Bering Sea, 752
projected impacts, 739, 740, 746, 759, 768, 1013	commercial harvest, 715
species, 487	harvest, 699
groundwater, 547	harvest by indigenous peoples, 652
and avalanches, 919	harvest of, 622, 628, 632
contribution to freshwater discharge, 225	population trends, 633
extraction and sea level, 231	projected impacts, 504, 510, 759
and frost heave, 916	hares ( <i>Lepus</i> spp.), 71, 268, 281, 300. see also individual species
and human health, 885	harp seal, 488
impacts of permafrost change, 45, 377	as cod predator, 737, 739, 741
impacts of projected change on fish, 402	commercial harvest, 699, 715, 735–736
impacts of projected change on freshwater ecosystems, 998	diet, 480
and landslides, 917–918	harvest by indigenous peoples, 652, 675
and methane emissions, 317	harvest of, 622–623, 625, 626, 628
systems, types of, 357–358	historic change, 663
grouse, 616–617, 654. see also hazel grouse; willow grouse; wood	impacts of capelin overharvesting, 624
grouse	impacts of past climate variability, 501, 739
growing degree-days, 801	in Northeast Atlantic, 696
for agriculture projections, 809–812	in northwest Atlantic, 732
crop requirements, 810	projected impacts, 504, 510, 740–741
impact of change on wetland vegetation, 392	Harpinia sp., 483
observed change, 1009	harvest regulation. see wildlife management
projected change, 391, 811–812	hay, 808–809
growing season length and timing, 245	hazel grouse (Tetrastes bonasia), 607
in boreal forests, 794	heating costs, 1004, 1007, 1013
and carbon balance, 274	heavy metals, 877-878, 947, 952, 982. see also cadmium; lead;
and contaminant transport, 890	mercury
impacts of projected change on carbon cycling, 391–392	analyses, 959
impacts of projected change on plants, 274	and human health, 890–892, 953–954, 964
impacts on agriculture, 807–809	impacts on biodiversity, 565
impacts on carbon balance, 317	in marine ecosystems, 624
observed change, 274	in reindeer meat, 979–980
projected change, 391	in traditional foods, 657
Grus americana. see whooping crane	Heinrich events, 49–50
guillemot, 392, 561, 626. see also black guillemot; pigeon	Helictotrichon krylovii, 258
guillemot; thick-billed murre	Hemilepidotus jordani. see yellow Irish lord
gulls, 358, 627–628, 629, 633, 634–635, 636. see also individual	Hemiptera, 553
species	hepatitis, 884, 887
Gulo gulo. see wolverine	herring ( <i>Clupea</i> spp.). see also individual species
Gwich'in, 81, 645, 680, 681	projected impacts, 504, 508, 660
observations of change, 660, 1014	herring gull ( <i>Larus argentatus</i> ), 624, 629–631, 630, 635
Gwich'in Council International, 13, 68–71, 602, 968	Heterocope spp., 387
Gymnocephalus cernuus. see ruffe	heterotrophic flagellates, 481–482
Gynaephora groenlandica, 557	hexachlorocyclohexanes, 891, 892, 952, 953, 964
gypsy moth (Lymantria dispar), 834	Hiatella arctica, 483
gyrfalcon (Falco rusticolus), 267, 556	Hiodon alosoides. see goldeye
gyrraicon (Faico Fasicolas), 207, 550	Hippoglossoides elassodon. see flathead sole
TT	
H	Hippoglossoides platessoides. see long rough dab
habitat	Hippoglossoides robustus. see northern flathead sole
availability and reproductive success, 418	Hippoglossus hippoglossus. see Atlantic halibut
classification and inventory needs, 582–583	Hippoglossus stenolepis. see Pacific halibut
defined, 543–544	HIRHAM regional climate model. see regional climate models
fragmentation, 605, 609, 616	hooded seal (Cystophora cristata), 488
impacts of fragmentation on biodiversity, 559, 566–567, 588,	as cod predator, 737, 739
1016	commercial harvest, 699, 735, 736
loss and reindeer husbandry, 976–977	food sources, 486
monitoring, 587	harvest by indigenous peoples, 652
projected change in extent, 570–571	harvest of, 622–623, 625, 626, 628
habitat protection. see protected areas; wildlife management	in northwest Atlantic, 732
haddock (Melanogrammus aeglefinus), 495, 500, 508, 696, 698,	projected impacts, 504, 510
704, 711, 712, 717, 740–741	Hormathia nodosa, 483

horned puffin (Fratercula corniculata), 626, 633, 635	impacts on Saami, 657
horse mackerel (Trachurus trachurus), 500	impacts on wildlife, 600
horsetail (Equisetum spp.), 363	impacts on wildlife habitat, 616
house mouse (Mus musculus), 605	projected impacts, 32, 188, 208, 223, 1001, 1008
Hudson Bay, Canada. see also marine Arctic	and protected areas, 604
indigenous observations of change, 68, 70, 661–662	and wildlife management, 638–639
productivity, 493	hydrogen, molecular, 312
projected change, 476–477	hydrogen sulfide, 544
variability of temperature and salinity, 469	hydroids, 580
Hudson Strait, Canada, 68, 70, 469. see also marine Arctic	hydroxyl radical, 155, 156, 317
human health	Hylocomium splendens, 257, 258, 265, 273, 275, 276, 277, 285,
and adaptation to climate change, 898–900	289–290, 571
alcohol abuse, 896	Hypogastrura tullbergi, 281, 556
causes of death, 868–869	Hypogastrura viatica, 282
and contaminants, 600, 615, 867, 890–892, 953–954, 964	Hypomesus olidus. see pond smelt
data for vulnerability assessments, 959	*
data on, 864	I
definition, 865, 893	ice, river and lake
and extreme events, 869–870	background, 220–221
and heat waves, 870	and freshwater ecosystems, 998
and housing infrastructure, 888–889 impacts of biodiversity change, 559	impacts of projected change, 223 impacts of projected change on freshwater ecosystems,
impacts of UV radiation, 953	997–998
importance of traditional foods, 653–654, 879, 956–957	impacts of projected change on indigenous subsistence
indicators, 898–900	activities, 668, 678, 683–684
indigenous observations of change, 651	impacts of projected change on infrastructure, 927
and industrial development, 608, 864	impacts of projected change on underwater UV, 419–420
infant mortality, 868	indigenous observations of change, 685, 993
life expectancy, 867–868	observed change, 31, 1009
links to biodiversity, 579	observed change in breakup timing, 220–221
multiple stresses, 874, 877–878, 893–895, 896–897, 947	observed change in freeze-up timing, 221–222
projected impacts, 1000, 1001–1002, 1004, 1009, 1011, 1014	projected change, 222, 923, 1012
projected impacts (direct), 869–878	research recommendations, 223
projected impacts (indirect), 878–892	and UV radiation, 996
research recommendations, 875, 877, 878, 897, 900–902, 982	ice grass (Phippsia algida), 258, 549
and sanitation infrastructure, 884, 886–888	ice jams, 23, 32, 221, 360, 380, 418–419, 885, 923
sexual abuse, 896	ice sheets. see glaciers and ice sheets
social, cultural, and mental health, 892-898	Iceland
and socio-cultural conditions, 865-868	agricultural impacts on, 799-800
suicide, 896	agriculture, 809
and temperature, 870–874	commercial fisheries, 709–731
and UV radiation, 874-878, 996	demographics, 868–869
and water supply infrastructure, 884–886	economic importance of fisheries, 721–722, 723–724
humpback whale (Megaptera novaeangliae), 489, 629, 715, 732,	forest area, 784
736, 753	forest tenure and management, 799-801
humpy shrimp (Pandalus goniurus), 749	historic deforestation, 799–800
hunting	invasive species, 605
adaptation options, 948	marine ecosystem description, 710–711
in Finnish forests, 796	observed change in glaciers and ice sheets, 206
impacts on biodiversity, 560–561	observed temperatures, 803
impacts on wildlife, 610, 613–615	overgrazing impacts, 555, 562
in Norwegian forests, 798	projected impacts on fisheries, 725–727
projected impacts, 558, 617, 622-623, 632, 634-636, 660,	projected impacts on forests, 800
1001	projected temperature change, 803
in Swedish forests, 797	reforestation, 800–801
technological change, 952	settlement, 250, 799–800
hunting, commercial	Iceland scallop (Chlamys islandica), 483
of marine mammals, 622–623	Iceland Sea, 493. see also marine Arctic
in Russia, 606–610	Icelus spiniger. see thorny sculpin
hunting, sport	icings. see aufeis
and boreal forests, 794	ide (Leuciscus idus), 406
economic benefits, 600, 626, 667, 674–675, 682	Igloolik, Nunavut, Canada
of marine mammals, 622	indigenous knowledge case study, 82–84
of polar bear, 657	indigenous observations of change, 68
of seabirds, 622	Illex illecebrosus. see squid
Hydrodates pelagicus, see European storm petrel	immunosuppression, 874–875, 996
Hydrodamalis gigas. see Steller sea cow	impacts assessment
hydroelectric generation	approaches, 8–10
and habitat loss, 977	impetigo, 884 impoundment, 359, 933
impacts of mild winter days, 23 impacts on indigenous peoples, 678	inconnu (Stenodus leucichthys), 654
impacts on indigenous peoples, 676 impacts on reindeer overgrazing, 601	meemia (bienouus teucteninys), 004
r	

indigenous knowledge and observations, 62-66, 646, 658	impacts of projected change in freshwater runoff, 229
academic engagement with, 64–65	impacts of projected change in permafrost, 215, 219
and adaptation to climate change, 92–93, 559, 669, 683–684,	impacts of projected change in sea ice, 195
955	impacts of projected change in sea level, 235–236
	1 1 3
of biodiversity change, 585	impacts of projected change in snow cover, 200
and biodiversity conservation, 558–559, 589	importance of traditional foods, 650–651, 653, 653–654,
of caribou, 960–961	656–657, 682–683, 686, 864–865, 879, 883,
case studies, 72–90	894–895, 953, 956–957, 1010
of change in caribou numbers, 611	land claim agreements, 13–14
of change in freshwater systems, 382	mixed economies, 656–657, 672–674, 682–683, 955, 956, 957,
of changing availability of traditional foods, 651	964, 983
of climate change, 66–90, 992–993	multiple stresses on, 64, 650–652, 657–658, 666, 676–678,
and co-management, 612, 613	678–680, 685, 686, 866–867, 948, 961, 964, 965,
"co-production of knowledge," 980	1001, 1014
defined, 9–10	observed impacts, 1010–1011
development of, 65	as part of food web, 583–584
and environmental impact assessment, 65	and permanent settlements, 664, 669, 678, 956, 1014
erosion of, 66, 90–91	prehistoric settlement, 13, 62, 64, 67, 246–247, 252–253,
of freeze-thaw events and caribou, 279	605–606
of frost sensitivity in plants, 276	projected impacts, 63-64, 68-71, 74-75, 76-77, 80-81, 82-84.
in Hudson Bay, 661–662	84–90, 90–93, 94–95, 1000–1001, 1000–1002, <i>1004</i> ,
ice predictability, change in, 80	1010–1011, 1013–1014, 1017
in impact studies, 947	projected impacts on livelihoods, 660–662
importance to survival, 64, 65	projected impacts on resource harvesting, 651, 658–660
and intellectual property, 65, 81	research recommendations, 686–687
1 1 2,7	
of landslides and subsidence, 377	resilience, 90–93, 650
links to scientific knowledge, 72, 77, 94	role in wildlife management, 600–602, 639–640, 644–648
of muskrat abundance, 386, 419	societal conditions and trends, 952, 955–957, 983
of "new" species of fish, 386	subsistence activities, 15, 246–247, 599, 605, 607, 608–610,
participatory research, 959–961, 962, 963, 982	610–611, 611–612, 614–615, 618, 622, 625–626,
of past climate change, 655–656	626–628, 628–631, 632–633, 634–636, 650–687,
of past climate change and polar bears, 674	747, 753, 764, 767, 865, 866, 868, 882–883, 894,
of past climate change and ringed seals, 674	947–948, 952–953, 961–962, 963, 964–965, 1000
and place names, 66	(see also hunting)
projected impacts, 416–417, 670	traditional food sharing, 654–655, 656
and reducing uncertainty, 333	vulnerability, 90–93
of reindeer/caribou, 267	vulnerability analysis, 948–949
research recommendations, 901	Indigenous People's Secretariat of the Arctic Council, 602
and resource management, 65	Indigirka Basin, 227
sea ice predictability, change in, 68, 76, 632, 667, 668, 669	individual transferable quotas (ITQs), 665
of spring snowmelt change, 376	industrial development. see also infrastructure; military operations;
and subsistence activities, 654	mineral extraction; oil and gas development;
and technological change, 868	transportation
of terrestrial ecosystems, 248	and human health, 864
and uncertainty, 329	impacts on ecosystems, 678
use of, 9–10, 65–66	impacts on abitats, 554–555, 566–567
of vegetation change, 271–272	impacts on indigenous peoples, 686
in vulnerability assessment, 948–949, 951–952, 957–961,	
	impacts on reindeer, 657
959–961, 980	impacts on wildlife, 609, 616, 620–621, 622, 639
weather predictability, change in, 68–69, 74, 78, 80, 81, 82–83,	projected impacts, 200, 219
84, 86, 87, 89, 94, 660, 662, 667, 667–668, 669, 670,	and wildlife management, 638–639
869–870, 883, 895, 963, 965, 993, 1013–1014	influenza. see avian influenza
and wildlife management, 634, 637, 638	infrastructure
indigenous peoples, 13–15, 62–64. see also specific peoples	definition and discussion, 926
acculturative stress, 893–896	engineering design for changing climate, 937–941
adaptation, 993, 1001	and habitat loss, 977
adaptation options, 91, 95, 662–666, 685	and human health, 884–890
adaptation to environmental variability, 62	impacts of projected change in permafrost, 215, 219, 660, 670,
adaptive strategies, 651, 955–957, 963	676, 678
barriers to adaptation, 663–665	impacts of projected change in snow cover, 200
coping strategies, 955–957	impacts of reduced sea ice, 76–77
cultural identity and wildlife, 654–655, 685	impacts of sea-level rise, 235–236
dependence on ecosystems, 4, 540	impacts of UV radiation, 996
economic impacts, 674–675	impacts on biodiversity, 555, 566–567, 567
fisheries, impacts on, 411–417	impacts on ecosystems, 247
and forest management, 792	indicators, 900
harvest and diet summary, 652–654	and natural hazards, 917–920
health status, 865–869, 870	observed damages, 935–936, 1011
historic adaptations, 662–663	on permafrost, 909–917
	1 ,
human rights and climate change, 64	projected impacts, 889, 926–935, 929–930, 939–940, 1000,
impacts of past climate change, 67 impacts of projected change in biodiversity, 558–559	1004, 1007, 1009, 1010, 1011, 1013
impacts of projected change in biodiversity, 558–559	research recommendations, 915, 941

risk analysis, 938	Inuktitut, 671
sanitation systems, 884, 886–888	Inupiaq, 599, 647, 674, 681
water supply, 884–886	and caribou co-management, 652
Innu, 680, <i>681</i>	caribou harvest, 680
insects	and government policy, 664
adaptation to arctic environment, 267	indigenous knowledge case study, 73–76
as bird prey, 552	and wildlife management, 632
1 37	<b>2</b> ,
and boreal forests, 785, 792, 830–835	Inupiat. see Inupiaq
impacts of projected change on plants, 278	Inuvialuit, 652, <i>681</i>
indigenous observations of change, 71, 78, 80, 86, 89, 287,	adaptation options, 666
<i>667</i> , <i>668</i> , <i>684–685</i> , <i>993</i>	adaptation strategies, 668–670
observed change, 1009	adaptation to observed change, 93
observed change in distribution, 1005	caribou harvest, 680
projected impacts, 570, 999, 1006, 1007, 1008, 1012	impacts case study, Sachs Harbour, 666–671
role in decomposition, 540	indigenous knowledge, 662
species numbers, 580	Inuit Observations of Climate Change project, 666–671
1	
wood-feeding, 552	observations of change, 585, 661, 961
insular vole (Microtus abbreviatus), 556–557	subsistence production, 653
Intergovernmental Panel on Climate Change (IPCC), 947	vulnerability case study, 963
on arctic climate change, 22–23, 31, 31, 37–38, 39, 42, 44	Inuvialuit Final Agreement, 91, 92, 613, 627, 665–666, 670
assessments, 6, 35	Inuvialuit Game Council, 632, 645, 666
on climate models, 102, 105	Inuvialuit-Inupiat Beluga Commission, 637
model choice, 109	invasive species, 555, 564, 575–577, 600. see also non-native
on precipitation increases, 186–187	species
projections of future global climate, 119	in Canada, 610
projections of future impacts, 961	definition, 605
on regional climate models, 132	indigenous observations of change, 80, 89
Special Report on Emissions Scenarios, 109	monitoring, 587
temperature projections, 120	and transportation corridors, 605
on thermohaline circulation, 125	and wildlife management, 605
on uncertainties in climate scenarios, 102	invertebrates
International Agreement on the Conservation of Polar Bears and	adaptations to cold, 556
their Habitat (IACPB), 623, 628, 632	diversity and latitude, 258
International Arctic Science Committee (IASC), 6	impacts of oil pollution, 565
International Convention for the Regulation of Whaling, 695, 965	monitoring recommendations, 587
International Council for the Exploration of the Sea (ICES), 698,	projected change in distribution, 569
699, 705	species numbers, 580
International North Pacific Fisheries Convention, 746–747	Ips typographus. see spruce engraver beetle
International Porcupine Caribou Board, 604	Igaluit, Nunavut, Canada
International Tundra Experiment (ITEX), 272–273, 585, 586	indigenous knowledge case study, 82–84
International Whaling Commission (IWC), 623, 626, 627, 629,	indigenous observations of change, 68, 70
633, 695, 705, 956, 965	Irish elk. see giant deer
	•
introduced species. see non-native species	island biogeography, 577–578
Inughuit, 652	isopods, 483
Inuit, 599, 602, 612, 652, 681. see also Inuvialuit	isoprene, 313
adaptation to climate change, 898	ivory gull (Pagophila eburnea)
adaptation to past climates, 672	projected impacts, 504, 512, 520, 999
and animal rights movement, 657	and sea ice, 480, 489, 490
barriers to adaptation, 664	status, 630
caribou harvest, 680	······································
demographics, 868	J
C 1 /	•
education and health, 892	jack pine budworm, 834
in Greenland, 866	jaegers. see skuas
health issues, 894–895, 896	James Bay, Canada, 68, 70, 662
historic adaptations, 662, 663–664	James Bay and Northern Quebec Agreement, 627, 665
impacts case study, Nunavut, 671–675	Janira maculosa, 483
impacts of past climate on, 67	jellyfish, 521, 758
importance of seal hunting, 650	Juniperus communis. see creeping juniper
importance of traditional foods, 654, 883	1 23 1
importance of whaling, 651	K
indigenous knowledge, 960–961	Kalaallit, 652
	,
observations of change, 82–84, 271, 279, 287, 615, 660,	Kallavesi Lake, Finland, 221
661–662, 961, 1014	Kangiqtugaapik. see Clyde River
oral histories, 655	Kara Sea, 493, 506
polar bear harvest, 627	Keewatin, 653
projected impacts, 1000	kelp, 480, 481
respiratory illness in, 870	Khanty-Mansiysky Okrug, Russia, 609
vulnerability to climate change, 866	killer whale (Orcinus orca), 489, 625, 629, 633, 713
and wildlife conservation, 626	king crab (Paralithodes camtschaticus), 704, 758
Inuit Circumpolar Conference, 13, 68–71, 602, 956, 968	king eider (Somateria spectabilis), 363, 573, 630, 633, 635, 653
Inuit Observations of Climate Change project, 666–671, 960	Kitikmeot Region, Canada, 69, 71, 660
Inuit Qauijimajatuqangit. see indigenous knowledge	

kittiwake ( <i>Rissa</i> spp.), 71, 624, 631, 653. see also individual	land use
species	and biodiversity, 1016
Kittlitz's murrelet, 635	change and carbon storage, 844–845
Kivalina, 660	impacts of change on biodiversity, 588
Kobresia spp., 262, 293	land use management, 621, 622
Koenigia islandica, 263	and biodiversity, 562–564 and wildlife management, 601, 602
Kola Peninsula, Russia, 604, 609, 965. see also Lovozero, Russia	
Kola Saami Reindeer Breeding Project, 666 Kolyma Basin, 227	landslides, 914 factors affecting slope stability, 917–918
	impacts of projected change in permafrost, 215, 913
Kolyma River, 229, 246 Koryak Autonomous Okrug, 559, 679	and infrastructure, 917
Kotzebue, Alaska	projected impacts, 917, 919–920, 1009, 1011, 1012
indigenous knowledge case study, 73–76	Lapland longspur (Calcarius lapponicus), 71, 258, 361
indigenous observations of change, 69	Laptev Sea, 493, 506, 518
Koyukon, 680	larch ( <i>Larix</i> spp.), 258, 262, 551, 784. see also individual species
krill, 482, 711–712	past distribution, 369
in Bering Sea, 754	projected impacts, 824
as prey, 486, 488, 489, 511	regeneration and temperature, 850
projected impacts, 758	in Russian boreal forest, 790
Kyoto Protocol, 119, 786, 791	treeline change, 847–848
,,,,	larch sawfly (Pristiphora erichsonii), 833–834
L	Larix cajanderi, see Cajander larch
Labrador. see Newfoundland and Labrador	Larix gmelinii. see Gmelin larch
Labrador Sea, 476–477, 493. see also marine Arctic	Larix laricina. see tamarack
Labrador tea ( <i>Ledum</i> spp.), 654	Larix sibirica. see Siberian larch
Lagenorhynchus acutus. see white-sided dolphin	Larix sukaczewii. see Sukachev larch
Lagenorhynchus albirostris. see white-beaked dolphin	Larus argentatus. see herring gull
Lagopus lagopus. see willow grouse	Larus canus. see mew gull
Lagopus mutus. see ptarmigan	Larus fuscus. see lesser black-backed gull
Lagopus mutus hyperboreus. see Svalbard ptarmigan	Larus glaucescens. see glaucous-winged gull
Lagotis minor, 258	Larus hyperboreus. see glaucous gull
lake chub (Couesius plumbeus), 405	Larus marinus. see great black-backed gull
lake cisco (Coregonus artedi), 398, 403	Larus ridibundus. see black-headed gull
Lake Saanajärvi, Finland, 382–383	Laysan albatross, 635
lake trout (Salvelinus namaycush), 564	Leach's storm petrel (Oceanodroma leucorhoa), 630
distribution, 395	lead mining. see mineral extraction
harvest by indigenous peoples, 652, 654, 667	lead pollution, 952, 954, 964
indigenous observations of change, 83	least auklet (Aethia pusilla), 489–490, 633, 635
projected impacts, 373, 378, 387, 403, 405, 407, 414, 435, 437,	least cisco (Coregonus sardinella)
1013	harvest by indigenous peoples, 667, 669
projected impacts on fisheries, 412	indigenous observations of change, 71, 386, 585, 667, 668
and water temperature, 398	least weasel (Mustela nivalis), 268, 301
lake whitefish (Coregonus clupeaformis)	Ledum spp. see Labrador tea
distribution, 397	lemmings ( <i>Lemmus</i> spp. and <i>Dicrostonyx</i> spp.). see also individual
projected impacts, 403, 408	species adaptation to arctic environment, 267–268
projected impacts on fisheries, 411 thermal preferences, 395, 398, 405	genetic variation, 285
lakes	observed change in population cycles, 279–280
area, 546	observed impacts, 279
and contaminants, 427–432, 437, 891	paleoecology, 249, 251
food web interactions, 564	population cycles, 268
freshwater ecosystems, 361–363	and predation, 302
indigenous observations of change, 88–90, 685	projected impacts, 281, 300, 998, 1005, 1006, 1017
inundation from sea-level rise, 383	role in ecosystems, 549–550
observed change in ice cover duration, 997	as "super-dominant" species, 260
perched, 31	trophic interactions, 298, 299–302
primary production, impacts on, 388–391, 435	Lemmus lemmus. see Norway lemming
projected change in carbon balance, 323–324, 393	Lemmus sibiricus. see brown lemming
projected change in ice cover duration, 570, 659, 997	Lemna spp. see duckweed
projected impacts, 381–384, 383, 412, 424, 434–435, 435, 437,	Lena Basin, 185–187, 187–188, 227, 228
998, 1012	Lena Delta, 610
and UV radiation, 424, 425-427, 435	Lena River, 222, 229, 358, 373, 381, 406–407
Laminaria spp., 481	lentic
Lamna ditropis. see salmon shark	definition of, 358
lamprey (Petromyzontidae), 396, 408, 576	Lepidopsetta bilineata. see rock sole
Lancaster Sound, 483	Lepomis macrochirus. see bluegill
land claims, 601, 612, 618, 627, 665. see also specific agreements	Lepus americanus. see snowshoe hare
land-fast ice, 30–31, 457. see also sea ice	Lepus arcticus. see Arctic hare
importance of, 31	Lepus timidus. see mountain hare
observed trends, 44	lesser black-backed gull (Larus fuscus), 629, 630
projected change, 472	lesser golden plover (Pluvialis dominica), 258
thickness measurements, 190	lesser scaup (Aythya affinis), 418

lesser snow goose (Anser caerulescens caerulescens), 280, 563	Mackenzie River
lesser white-fronted goose (Anser erythropus), 288, 574, 1006	and carbon transport, 518
Leuciscus idus. see ide	freshwater ecosystems, 358
Leuciscus leuciscus baicalensis. see common dace	observed change in freeze-up date, 221
lichens, 549. see also microorganisms	observed change in ice duration, 423
adaptations to arctic environment, 269–270	projected impacts, 229, 373, 375–376, 381
arboreal, 797	mackerel. see also individual species
canopy, 552	in northwest Atlantic, 732
diversity, 1016	past distribution change, 500
genetic response to change, 285	projected impacts, 508, 700, 721, 740, 1007
impacts of overgrazing, 563	macroalgae
importance to reindeer/caribou, 660–661, 680	biomass, 480
indigenous observations of change, 272	in marine ecosystems, 481
knowledge of arctic species, 540	Maldane sarsi, 483
projected impacts, 967, 997, 998, 1006, 1017	mallard (Anas platyrhynchos), 287, 669
species numbers, 580	Mallotus villosus. see capelin
vulnerability to warming, 273	mammals. see also wildlife; individual species
light-bellied brent goose (Branta bernicla hrota), 574	diversity, 1016
lightning	impacts of change on human health, 879–892
• •	indigenous observations of change, 993
and wildfires, 836–839	
likelihood	species numbers, 580
terminology (lexicon), 6–7, 993, 994	mammals, aquatic. see also wildlife; individual species
Limacina helicina, 482	projected impacts, 417–419, 436
Limanda aspera. see yellowfin sole	mammals, marine, 487–490. see also wildlife; individual species
Limnodromus scolopaceus. see long-billed dowitcher	adaptations to cold, 556–557
Limosa lapponica. see bar-tailed godwit	and animal rights movement, 965
Limosa limosa. see black-tailed godwit	in Bering Sea, 746
Lincoln Sea, 493	climate impacts on, 496–497
ling (Molva molva), 500, 711, 717	and commercial fisheries, 708
lingonberry (Vaccinium vitis-idaea), 273, 275, 276, 276, 277	commercial harvest, 705, 752–753
impacts of UV radiation, 571	and contaminants, 600, 602
indigenous observations of change, 271–272	harvest in Fennoscandia, 628–631
Lithodes aequispina, 483	harvest in Russia, 609
little auk (Alle alle), 489–490, 712	harvest of, 622–634, 952
harvest of, 627, 631	and human health, 896–897
in northwest Atlantic, 732	impacts of changing sea ice on harvests, 195
projected impacts, 512, 520	impacts of commercial harvesting, 559
and sea ice, 480	impacts of oil and gas development, 624
status, <i>630</i>	impacts of past climate variability, 504, 699–700, 716–719
little bunting (Emberiza pusilla), 258	impacts of projected change on subsistence activities, 667
Little Climatic Optimum. see paleoclimate	impacts of reduced sea ice, 952
Little Diomede, 660	importance to indigenous peoples, 599–600, 622, 632, 634
Little Ice Age. see paleoclimate	indigenous harvesting, 747, 1000
little stint (Calidris minuta), 258, 288	indigenous observations of change, 662, 993
Littorina saxatilis, 483	inventory needs, 580
liverworts, 540, 580	management and conservation, 622-636, 639, 695, 747
livestock, 807-809, 977	mercury levels in, 566, 964
lodgepole pine (Pinus contorta), 800	observed change, 1009, 1010
long-billed dowitcher (Limnodromus scolopaceus), 288	overharvesting, 487, 601, 606
long rough dab (Hippoglossoides platessoides), 484, 487, 732, 737	projected change in distribution, 546, 588, 700
long-tailed duck (Clangula hyemalis), 71, 86, 363, 629–630, 630	projected impacts, 504, 509–511, 545–546, 604, 658–659, 768.
long-tailed jaeger (Stercorarius longicaudus), 361	880, 1010, 1013
Lophelia pertusa, 561	research recommendations, 631
Lota lota. see burbot	threats to migratory species, 581
lotic	mammals, terrestrial. see also wildlife; individual species
definition of, 358	adaptations to cold, 556–557
Lovozero, Russia	diversity and latitude, 258
indigenous knowledge case study, 87–90	indigenous observations of change, 662
indigenous observations of change, 87–90	number of species, 556
Lund-Potsdam-Jena model, 323	overharvesting, 601, 606
lupine ( <i>Lupinus</i> spp.), 605	projected impacts, 560, 658–659
Lupinus spp. see lupine	mammoth ( <i>Mammuthis primigenius</i> ), 246–247, 250, 252, 584
Lutz spruce, 831	Mammoth steppe. see steppe-tundra
Luzula confusa, 247	Mammuthis primigenius. see mammoth
Lymantria dispar. see gypsy moth	marbled murrelet, 635
Lymnaea elodes, 564	marine Arctic. see also Arctic Ocean
lynx (Lynx lynx), 601, 619, 977	circulation, 454–456, 458, 460–465, 710–711
	circulation, importance to ecosystems, 490
M	circulation, observed change, 467–469
Mackenzie Basin, 185–187, 187–188, 227, 1000, 1011	circulation, projected change, 472–473, 476
Mackenzie Basin Impact Study, 947, 960	circulation, uncertainties, 521
Mackenzie Delta, 80, 547	circulation, uncertainties, 321
MIGUNOLIZIU DUIG, OU, JT/	Circulation and fish distribution, 470

1 1 1 1 1 17	
circulation and sea ice, 471	marine management
circulation and wind, 461	and biodiversity, 562–564
currents, 464	and wildlife management, 601
currents and ecosystems, 490	marsh. see wetlands
feedbacks to global climate, 459	marten, 619
freshwater budget, 460	Martes zibellina. see Sable
freshwater storage, 459	mattak, 652, 895
heat balance, 459	Medieval Warm Period. see paleoclimate
links to global climate, 458–465, 459, 473, 519–520	megafauna
mixed layer depth, 460–461	extinction hypotheses, 584
ocean fronts, 477, 483, 521	Megaloceros giganteus. see giant deer
physical characteristics, 454–478	Megaptera novaeangliae. see humpback whale
prehistoric change, 466	Melanitta fusca. see velvet scoter
projected change, 469–478, 473–475, 476	Melanitta nigra. see black scoter
research recommendations, 520–522	Melanogrammus aeglefinus. see haddock
stratification, 460	meningitis, 884
variability, 466–469	Menyanthes spp., 317
water budget, 465	mercury, 952, 954, 982
marine ecosystems	in aquatic ecosystems, 432–433, 565
area, 544	in arctic ecosystems, 566
benthos, 482–483	depletion events (MDEs), 424, 432, 891, 955
of Bering Sea, 747, 754–755, 755	global emissions of, 432
biodiversity conservation, 544–546	in human diets, 964
biodiversity monitoring, 586	impacts of change in UV radiation, 424
carbon biomass and productivity, 479	impacts of climate change on, 432–433
and carbon cycle, 518	impacts on biodiversity, 565
characteristics, 1017	sources of, 615
climate impacts on, 490–491	and traditional foods, 883
and contaminants, 610	transport pathways, 891
described, 12, 479–490	Mergus merganser. see common merganser
distribution of species, 499	Mergus serrator. see red-breasted merganser
food-web energy transfer, 494	meridional overturning circulation. <i>see</i> thermohaline circulation
food webs, 479–480	Metarhizium, 270, 283
food webs and UV radiation, 514	meteorological stations, 910
freshwater impacts on nutrient supply, 544	methane
and human health, 882	as bacterial energy source, 544
ice cover, importance of, 12	contribution to carbon balance, 317–318
of Iceland and Greenland, 710–712	global warming potential, 318
impacts of aquaculture, 564	and hydrogen oxide (hydroxyl) radical, 156, 317
impacts of ballast water discharge, 576	and microorganisms, 260
impacts of overharvesting, 624, 658, 659 impacts of past climate variability, 497–498, 500, 503, 658	observed flux change, 1005 and ozone chemistry, 155–156
impacts of past chinate variability, 497–498, 300, 303, 038 impacts of projected change in glaciers and ice sheets, 208	and permafrost degradation, 319, 459
impacts of projected change in graciers and ice sneets, 208	projected change in fluxes, 998, 999
impacts of projected change in permanost, 219	as sink for reactive chlorine, 155–156
impacts of projected change in sea level, 235	soil processes affecting, 313
impacts of projected change in sea level, 233	soil sources and sinks, 312
impacts of projected precipitation change, 188	and stratospheric temperature, 155
impacts of UV radiation, 947, 996	as water vapor source, 155
impacts of UV radiation, 547, 550	methane emissions
limiting nutrients, 491	environmental controls, 312–313
Newfoundland and Labrador, 731	impacts of permafrost degradation, 34, 219, 391–393
in Northeast Atlantic, 696	impacts of projected change in active-layer depth, 34
of northwest Atlantic, 732–733	impacts of projected change in lakes, 393
observed change, 659	from marine methane hydrates, 34
and ocean temperature, 659	observed change, 1006
primary production, 479–480, 483, 491–493, 520	oceanic, 519
primary production and UV radiation, 514–515	and plant species composition, 310–311, 317
productivity, 12	projected impacts, 1008
projected impacts, 504, 504–512, 507, 544–546, 560, 569, 570,	and projected vegetation change, 323
694, 758, 968–969, 999, 1008	recent change in terrestrial ecosystems, 317
protected areas, 577	and sedges, 294
research recommendations, 520–522	and soil temperature, 317
in Russian Arctic, 623–624	from terrestrial ecosystems, 309, 317–318
and salinity, 659	and UV radiation, 310
sea ice communities, 480	from wetlands, 364
secondary production, 493–494	methane hydrates, 519, 999, 1014, 1015
sensitivity to climate change, 521	marine, impacts of temperature increase, 34
species diversity, 12	potential for release, 219
trophic levels defined, 479	in subsea permafrost, 217
uncertainties in projecting change, 693–694	terrestrial, 34
	Methylocansa 260

Mathylacolla 260	species numbers, 580
Methylocella, 260 Metis, 681	mitigation, 1020
Metridia spp., 482	definition, 951
mew gull (Larus canus), 630	and human health, 865
microalgae, 481	models. see climate models
Microgadus proximus. see Pacific tomcod	Mola mola, see ocean sunfish
microheterotrophs, 481–482. see also bacteria	mollusks
Micromesistius poutassou. see blue whiting	commercially harvested species, 483
microorganisms. see also bacteria	role in decomposition, 540
adaptations to arctic environment, 269–271	species numbers, 580
community response to experimental manipulation, 297–298	Molva molva. see ling
difficulty studying, 330	Monodon monoceros. see narwhal
diversity, 260–262	Montreal Protocol, 169, 174, 176, 995
genetic response to change, 286	moose (Alces alces), 301, 897
projected change in distribution, 291	harvest by indigenous peoples, 652-653, 654
projected impacts, 282–284	harvest in Fennoscandia, 616–617
in sea ice, 480	harvest in Russia, 607, 609
trophic interactions, 303	indigenous observations of change, 70–71, 73, 80, 87, 879
and UV radiation, 270	as invasive species, 605, 610
winter respiration, 269–270	observed change, 1005
Micropterus dolomieu. see smallmouth bass	population monitoring, 620
Microtus abbreviatus. see insular vole	population status in Alaska, 619
Microtus gregalis. see narrow-headed vole	projected impacts, 660, 1010, 1011
Microtus middendorffi. see Middendorf's vole	Morus bassanus. see northern gannet
Microtus oeconomus. see tundra vole	mosquito, 364, 1017
Microtus rossiaemeridionalis. see sibling vole	indigenous observations of change, 71, 86, 89
Microtus spp. see voles	moss campion (Silene acaulis), 285, 582
Middendorf's vole (Microtus middendorffi), 258	mosses, 549, 552, 797
midge, 1017. see also chironomids migratory species, 1015, 1017	diversity, 1016 genetic response to change, 285
impacts of human development, 555	knowledge of arctic species, 540
indigenous observations of change, 585, 685	projected impacts, 275, 997, 998, 1006, 1017
projected change in routes, 418, 588	species numbers, 580
projected change in fouces, 716, 566	true mosses, 293
projected impacts, 570, 658–659, 660	vulnerability to warming, 273
wildlife management challenges, 599, 601–602, 639–640	Motacilla alba. see white wagtail
Milankovitch, 47	moths, 552, 571
military operations	mountain avens (Dryas integrifolia), 257, 258, 275, 285, 293, 582
impacts of projected change in sea ice, 195	mountain birch (Betula pubescens ssp. czerepanovii), 281, 292,
impacts on habitat, 616, 977	295, 304, 834, 967, 1005–1006
impacts on habitat, 616, 977 impacts on indigenous peoples, 679	295, 304, 834, 967, 1005–1006 mountain crowberry (Empetrum hermaphroditum), 276, 276, 277
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008,	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666,
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961,
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961,
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715 marine biomass, 479 marine productivity, 479 in northeast Atlantic, 696	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620 Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020 on seabirds, 545 vulnerability assessment, 963, 965, 981 on wildlife, 602, 615
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on reindeer, 657 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715 marine biomass, 479 marine productivity, 479 in northeast Atlantic, 696 in northwest Atlantic, 732	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020 on seabirds, 545 vulnerability assessment, 963, 965, 981 on wildlife, 602, 615 Munna sp., 483
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on vildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715 marine biomass, 479 marine productivity, 479 in northeast Atlantic, 696 in northwest Atlantic, 732 population, 736	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020 on seabirds, 545 vulnerability assessment, 963, 965, 981 on wildlife, 602, 615 Munna sp., 483 murres (Uria spp.). see also individual species
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on vildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715 marine biomass, 479 marine productivity, 479 in northeast Atlantic, 696 in northwest Atlantic, 732 population, 736 projected impacts, 510	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020 on seabirds, 545 vulnerability assessment, 963, 965, 981 on wildlife, 602, 615  Munna sp., 483 murres (Uria spp.). see also individual species harvest of, 626, 633
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on wildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715 marine biomass, 479 marine productivity, 479 in northeast Atlantic, 696 in northwest Atlantic, 732 population, 736 projected impacts, 510 mire. see wetlands	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020 on seabirds, 545 vulnerability assessment, 963, 965, 981 on wildlife, 602, 615  Munna sp., 483 murres (Uria spp.). see also individual species harvest of, 626, 633 indigenous observations of change, 71
impacts on indigenous peoples, 679 mineral extraction, 16, 990 and biodiversity, 566 and ecosystems, 247 impacts of projected change in river and lake ice, 229 impacts of projected change in sea ice, 195 impacts on arctic residents, 1000 impacts on indigenous peoples, 678–680 impacts on vildlife, 600, 601, 602, 609, 616, 619, 620–621 projected impacts, 519, 926–927, 937, 1001, 1002, 1004, 1008, 1013 and protected areas, 604, 604 role in mixed economies, 656 in Russia, 606 and wildlife management, 638–639 mink (Mustela vison), 619, 1005 minke whale (Balaenoptera acutorostrata), 489, 615 commercial harvest, 699, 705 harvest by indigenous peoples, 652, 965 harvest of, 622, 628, 629, 633 historic harvesting, 715 marine biomass, 479 marine productivity, 479 in northeast Atlantic, 696 in northwest Atlantic, 732 population, 736 projected impacts, 510	mountain crowberry (Empetrum hermaphroditum), 276, 276, 277 mountain hare (Lepus timidus), 280, 552 mountain sheep population monitoring, 620  Mucor hiemalis, 298 mule deer (Odocoileus hemionus), 78 multiple stresses on Arctic ecosystems, 599, 605, 636 on arctic residents, 946–949, 950–952, 1000 on boreal forests, 566, 797 on coastal communities, 771 on ecosystems, 247–248 on fisheries, 706–707, 708 on freshwater ecosystems, 548, 998 and human health, 874, 877–878, 893–895, 896–897 on indigenous peoples, 64, 94–95, 650–652, 657–658, 666, 670, 676–678, 678–680, 685, 686, 866–867, 961, 964, 1001, 1014 on marine ecosystems, 509, 515, 999 on marine mammals, 545 on reindeer husbandry, 970–971, 980 research recommendations, 1019, 1020 on seabirds, 545 vulnerability assessment, 963, 965, 981 on wildlife, 602, 615  Munna sp., 483 murres (Uria spp.). see also individual species harvest of, 626, 633

musculoskeletal diseases, 873	Nelson Lagoon, Alaska
mushrooms, 792, 796, 797	indigenous knowledge case study, 76–77
muskeg, 32	nematodes, 297, 303, 480, 483, 554
muskox (Ovibos moschatus), 252, 570, 1016	role in decomposition, 540
adaptation to arctic environment, 267	species numbers, 580
and caribou, 661	Nenets, 652
commercial harvest, 667	impacts case study, 675–678
farming, 808, 813	population, 679
genetic diversity, 582	Nenetsky Okrug, Russia, 609
harvest by indigenous peoples, 652–653, 667, 669	Neodiprion sertifer. see European pine sawfly
harvest management in Canada, 612	net ecosystem production
impacts of ice crusting, 279, 611–612	projected change, 326
impacts of observed change, 615	of terrestrial ecosystems, 306–309
impacts of warming on lungworm, 302	net primary production
impacts on vegetation, 300	impacts of experimental treatments, 296
indigenous observations of change, 71, 271, 585	projected change, 321, 326
observed change, 675, 1005	spatial variability, 306
observed change on Banks Island, 666–667	of terrestrial ecosystems, 306, 307–309
overharvesting in Canada, 611, 670	uncertainty in measuring, 330
· · · · · · · · · · · · · · · · · · ·	
paleoecology, 250	Netsilik, 672, 674
population cycles, 268	Newcastle disease, 880
population increase in Yakutia, 609	Newfoundland and Labrador, 653, 731–746, 805
population monitoring, 620	nickel. see mineral extraction
population status in Alaska, 619	ninespine stickleback ( <i>Pungitius pungitius</i> ), 385, 397
* *	
and predation, 301–302	ningiqtuk, 671, 672–674
projected impacts, 1012	nitric oxide, 313
threats from hunting, 613–614	nitrogen
muskox lungworm, 302	in boreal forest soils, 795, 797, 798, 820–821, 843
muskrat (Ondatra zibethicus), 363	cycle and terrestrial ecosystems, 305–311
//	
harvest by indigenous peoples, 653, 654	deposition and biodiversity, 564–565
importance of estuaries, 360	deposition and boreal forests, 845–846
projected impacts, 386, 418–419	deposition and treeline advance, 851
and tularemia, 881	deposition and vegetation distribution, 568
mussel, 745. see also blue mussel	nitrous oxide
Mustela erminea. see ermine	and ozone chemistry, 155–156
Mustela nivalis. see least weasel	soil processes affecting, 313, 314
Mustela vison. see mink	soil sources and sinks, 312
mustelids, 300–301	and stratospheric temperature, 155
Mya arenaria. see clams	nodding saxifrage (Saxifraga cernua), 285
·	
Mya truncata. see clams	Noemacheilus barbatulus. see stone loach
Myoxocephalus quadricornis. see fourhorn sculpin	non-Hodgkins lymphoma, 876–877
Mysis relicta. see opossum shrimp	non-native species, 247, 571, 999, 1006. see also invasive species
Mytilus edulis. see blue mussel	in boreal forests, 804, 834
<b>,</b>	and climate change, 576
N.	
N	in Greenland, 584
naleds. see aufeis	and human health, 879, 880
nanoflagellates, 506	and hybridization, 573
narrow-headed vole (Microtus gregalis), 258	impacts of introduced species, 563
narwhal (Monodon monoceros), 489	impacts on biodiversity, 542, 545, 547, 548, 575–577
· · · · · · · · · · · · · · · · · · ·	
food sources, 486	indigenous observations of, 70–71, 77, 615, 660, 667, 993, 993
harvest by indigenous peoples, 652, 675	projected change, 281, 509, 555
harvest management in Canada, 612, 627	and wetland drainage, 562
harvest of, 629	Nordic Seas, 475–476. see also marine Arctic
historic change, 663	Norman Wells Pipeline, 932
importance of sea ice, 659	North Atlantic Deep Water, 461, 464, 473
migration, 546	North Atlantic Marine Mammal Commission (NAMMC), 623,
projected impacts, 504, 510–511, 999	666, 695, 705, 956
Naskapi, 680	North Atlantic Ocean, 468–469. see also marine Arctic
natsiq. see ringed seal	North Atlantic Oscillation (NAO), 24–25, 974, 990, 1018. see also
natural hazards. see also avalanches; floods; landslides	
	Arctic Oscillation
and infrastructure, 917–920	and cooling in northwest Atlantic, 732–733
model projections, 927–928	and copepod abundance, 497
natural resources, 15–16. see also fisheries; mineral extraction; oil	and fish distribution, 500
and gas development	impacts on Arctic climate, 10
navigation. see also Northern Sea Route; Northwest Passage	•
	impacts on ocean temperature, 572
impacts of projected change in freshwater runoff, 229	and the Little Ice Age, 53
impacts of projected change in glaciers and ice sheets, 208	and ocean circulation, 465, 467, 469
impacts of projected change in precipitation, 195	and ozone depletion, 157
impacts of projected change in river ice, 223	and pine growth, 829
impacts of projected change in sea ice, 195	projected impacts, 119, 134–135, 470, 740
projected impacts, 566, 997, 1008	and river discharge, 226
NCEP/NCAR reanalysis, 111–117	and sea ice duration, 469

North East Atlantic Fisheries Commission (NEAFC), 694, 706	economic importance of fisheries, 700–704
North Pacific. see also Bering Sea; marine Arctic	fisheries management, 704–706
commercial fisheries, 746–770	forest area, 784
North Pacific climate regime shift, 633. see also Bering Sea	forest tenure and management, 798–799
North Pacific Oscillation, 572	protected areas, 541
North Pacific right whale (Eubalaena japonica), 753	sanitation systems, 884
North Water Polynya, 30, 477, 493	vulnerability assessment, 968–969
Northeast Passage. see Northern Sea Route	Norway lemming ( <i>Lemmus lemmus</i> ), 268, 300, 301, 549
Northeast Water Polynya, 493	Norway rat (Rattus norvegicus), 605
northern anchovy (Engraulis mordax)	Norway spruce ( <i>Picea abies</i> ), 252, 292
impacts of UV radiation, 513	in Finland, 795
Northern Annular Mode. see Arctic Oscillation (AO)	projected change in distribution, 568, 584
northern bluefin tuna. see bluefin tuna northern bluegrass (Poa abbreviata), 258	projected impacts, 845 soil warming experiment, 820–821
northern flathead sole (Hippoglossoides robustus), 751	Norwegian Sea
northern fulmar (Fulmaris glacialis), 361, 489	commercial fisheries, 695–709
harvest of, 626, 629–630	ecosystem description, 696
in northwest Atlantic, 732	"Herring hole," 695
status, 630, 635	productivity, 493
northern fur seal (Callorhinus ursinus), 77, 489	projected change in fish distribution, 508
commercial harvest, 752–753	Notidanus griseus. see bluntnose sixgill shark
harvest of, 623, 625	Notropis atherinoides. see emerald shiner
historic harvesting, 761	Novaya Zemlya
impacts of POPs, 566	observed change in glaciers and ice sheets, 206
indigenous harvesting, 753	Numenius borealis. see Eskimo curlew
management regimes, 746	Nunamiut, 680
observed change, 1009	nunataks, 249, 550–551
population trends, 633	Nunavummiut, 671, 673
northern gannet (Morus bassanus), 629, 630	Nunavut
northern goshawk, 89	demographics, 868
northern lapwing (Vanellus vanellus), 287	impacts on indigenous peoples case study, 671–675
northern phalarope (Phalaropus lobatus), 489, 630	indigenous knowledge case study, 82–84
northern pike (Esox lucius)	Nunavut Land Claims Agreement, 627, 665
distribution, 395, 396–397	Nuorgam, Finland
fisheries, 411	indigenous knowledge case study, 84–87
harvest by indigenous peoples, 652	nutrient cycling
indigenous observations of change, 70–71	impacts of projected change on biodiversity, 589
northward colonization, 406	Nyctea scandiaca. see snowy owl
projected impacts, 1013	
and water temperature, 398, 405, 407-408	O
northern redbelly dace (Chrosomus eos), 396	oats, 807–808, 810
Northern Sea Route, 934–935, 1002, 1004	Ob Basin, 185–187, 187–188, 227, 552, 608
and coastal erosion, 519	Ob River
impacts of increased traffic, 555, 678, 962	and carbon transport, 518
and natural resource development, 937, 1008	contaminant loading of, 381
projected change, 519, 922–923, 924–926, 999	dissolved oxygen in, 379
northern shrimp (Pandalus borealis), 483	freshwater discharge, 547
in Central North Atlantic, 711–712	freshwater ecosystems, 358
commercial fishery, 697, 713, 714, 715, 722–723, 734–735,	impacts of reduced ice on reindeer husbandry, 678
738–739, 749	projected impacts, 229, 373, 381
commercial harvest, 652	obesity, 883, 897, 1000
harvest by indigenous peoples, 652	ocean currents
impacts of past climate variability, 719, 738–739	and contaminant transport, 891
importance to Greenland economy, 657	indigenous observations of change, 661
in northeast Atlantic, 696	Ocean Model Intercomparison Project (OMIP), 109
in northwest Atlantic, 732–733	ocean sunfish (Mola mola), 500
projected impacts, 720, 728, 740–741, 1007, 1013	Oceanodroma leucorhoa. see Leach's storm petrel
northern wheatear ( <i>Oenanthe oenanthe</i> ), 258	Ochejohka, Finland
Northwest Atlantic Fisheries Organization (NAFO), 694, 731	indigenous knowledge case study, 84–87  Odobenus rosmarus, see walrus
Northwest Passage, 1002, 1004 impacts of increased traffic, 555, 1013	Oenanthe oenanthe. see northern wheatear
and natural resource development, 937	official burnet (Sanguisorba officinalis), 258
navigation and sea ice, 196	oil and gas development, 15–16, 990
projected change, 519, 922–923, 924, 934, 1012	and biodiversity, 566
Northwest Territories	and ecosystems, 247
agriculture, 807–808	and habitat damage, 609
demographics, 868–869	impacts of projected change in permafrost, 219
indigenous subsistence production, 653	impacts of projected change in permanost, 219
Norway	impacts or projected change in river and take ice, 229
agriculture, 808–809	impacts on caribou, 621, 646
commercial fisheries, 695–709	impacts on indigenous peoples, 678–680, 685
4demographics, 868	impacts on marine ecosystems, 624

impacts on reindeer, 657	and UV radiation, 875
impacts on reindeer herding, 962	
impacts on wildlife, 600, 601, 602, 619, 620-621	P
impacts on Yamal Nenets, 676–678	Paamiut, Greenland, 663–664
importance to Alaskan economy, 657	Pacific cod (Gadus macrocephalus), 486, 495
observed damage to pipelines, 936	commercial fisheries, 750–751, 762, 766–767
projected impacts, 519, 926, 927, 931–933, 934, 936–937,	distribution, 484
<i>1001</i> , 1002, <i>1004</i> , 1010, 1013	historic harvesting, 761–762
and protected areas, 604, 604, 793	impacts of past climate variability, 503, 503, 755–756, 766
role in mixed economies, 656	projected change in distribution, 545
and wildlife management, 638–639	projected impacts, 507, 758, 759–760, 768
oil pollution	Pacific Decadal Oscillation (PDO), 26, 990
impacts on biodiversity, 565–566	impacts on arctic climate, 10
oilseed, 809	impacts on Pacific salmon, 757
Okhotsk Sea, 493	and marine ecosystems, 26
Old Crow, Yukon Territory, 654, 680–685	and ocean temperature, 469
oldsquaw. see long-tailed duck	reconstruction with tree rings, 819
oligochaetes, 483	and sea ice, 469
Oncorhynchus gorbuscha. see pink salmon	and wildfires, 836
, .	
Oncorhynchus keta. see chum salmon	Pacific golden plover ( <i>Pluvialis fulva</i> ), 258
Oncorhynchus kisutch. see coho salmon	Pacific halibut (Hippoglossus stenolepis)
Oncorhynchus mykiss. see rainbow trout	bycatch, 751
Oncorhynchus nerka. see sockeye salmon	commercial fisheries, 762, 764
Oncorhynchus tshawytscha. see chinook salmon	historic harvesting, 761
Ondatra zibethicus. see muskrat	impacts of past climate variability, 766
Onisimus spp., 482	indigenous harvesting, 747
Onychiurus spp., 282	management regimes, 746, 765
Operophtera brumata. see winter moth	projected impacts, 758, 759
Ophiopholis aculeata, 483	Pacific herring (Clupea pallasi), 485
opossum shrimp (Mysis relicta), 563	commercial fisheries, 748
oral histories. see indigenous knowledge	distribution, 484
Orcinus orca. see killer whale	historic harvesting, 761–762
organochlorine compounds, 624, 657. see also contaminants	impacts of past climate variability, 503, 503, 755–756, 766
Orok people, 679	indigenous harvesting, 747
Orthothecium chryseon, 258	management regimes, 765
Osmerus mordax. see smelt	projected change in distribution, 545
osprey, 71	projected impacts, 507, 758, 759–760, 768, 770–771, 1010
Ovibos moschatus. see muskox	Pacific loon (Gavia pacifica), 633, 635
Ovis canadensis dalli. see Dall sheep	Pacific mackerel (Scomber japonicus), 513
owls (Asio, Nyctea spp.), 268, 279, 291. see also short-eared owl;	Pacific ocean perch, 762, 766
snowy owl	Pacific salmon ( <i>Onchorhynchus</i> spp.). see also individual species
Oxycoccux spp. see cranberry	commercial fisheries, 748, 752, 762–763, 763, 764, 767
Oxyria digyna. see Arctic sorrel	and contaminant transport, 431, 892
oysters, 880	historic harvesting, 761
ozone, total column	impacts of past climate variability, 757, 766, 879
factors affecting, 154–157	indigenous harvesting, 747
feedback to climate change, 171	indigenous observations of change, 71, 386, 668
future recovery, 169, 174	management regimes, 746, 765
monitoring instruments, 158	projected distribution change, 410
observations and projections compared, 172, 173	projected impacts, 758, 759, 760, 768
observed change, 157–159, 248	projected impacts on fisheries, 415
projected change, 169–174	Pacific tomcod (Microgadus proximus), 486
research recommendations, 175–176	Pagophila eburnea. see ivory gull
seasonal variation, 996	paleoclimate, 46–54, 466, 1018
transport processes, 156–157	in Central Eurasia, 814–817
uncertainties in projections, 173	Holocene, 50-52, 365-371
and UV radiation, 153, 158, 160, 161–162, 166	last glacial/interglacial, 49–50
ozone, tropospheric	Last Glacial Maximum, 48–49, 249
and UV radiation, 153	last glaciation (Wisconsinan), 48–49
ozone depletion, 994–995	last interglacial (Eemian), 48
Antarctic ozone hole, 154, 157, 170	last millennium, 52–54
and biodiversity, 1016	Little Climatic Optimum, 671–672
impacts of climate change, 153, 154, 170–171, 419	Little Ice Age, 53–54, 67, 250, 466, 655–656, 671–672, 816,
•	• · · · · · · · · · · · · · · · · · · ·
and mercury depletion events, 891	820, 990 Medieval Warm Pariod 52 52 466 674 814 816 817 820
and meteorological variability, 156–157	Medieval Warm Period, 52–53, 466, 674, 814, 816–817, 820,
"mini-holes," 157	848, 990
modeling, 142–143, 171–174	Neo-Atlantic Optimum, 663
observed, 5, 154, 874, 996	in northwestern Europe, 819–820
ozone-depleting substances, 154, 176, 995	pre-Quaternary Period, 46–47
and polar stratospheric clouds, 419	proxy records of, 67, 365–366
and the polar vortex, 419	Quaternary Period, 47
projected, 996	rapid climate change, 23, 49–50, 51, 54, 125, 249–250

reconstruction with tree rings, 814–820	permafrost, terrestrial. <i>see also</i> thermokarst
and salmon fluctuations, 658	active layer defined, 209–210
Scandic Period, 663	active layer depth, 245
temperature variability, 23	active layer detachment, 549, 917
on Yamal Peninsula, 847–848	and arctic wetlands, 45–46
Younger Dryas, 50, 249, 466 Paleoclimate Modeling Intercomparison Project (PMIP), 109	area of, 45
- · · · · · · · · · · · · · · · · · · ·	background, 209–210 in boreal forests, 794
Pallas's cormorant, 626	
paludification, 320, 550, 678, 835, 851 pandalid shrimp ( <i>Pandalus jordani</i> ), 749, 759, 768	and carbon balance, 319, 841 carbon storage, 305, 518, 1014
Pandalopsis dispar. see deepwater prawn	carbon storage, 303, 316, 1014
Pandalus borealis. see northern shrimp	continuous, defined, 209
Pandalus goniurus. see humpy shrimp	degradation, impacts on buildings, 889, 929–930, 935, 1009
Pandalus jordani. see pandalid shrimp	degradation, impacts on carbon balance, 34, 391–393, 459
Papaver polare. see Arctic poppy	degradation, impacts on DOC, 391–393
Papaver spp., 293	degradation, impacts on freshwater ecosystems, 434–435,
paper birch (Betula papyrifera), 576, 792, 828, 834	997–998
Paragorgia arborea. see red gorgonian	degradation, impacts on infrastructure, 926–928, 939–940,
parakeet auklet (Aethia psittacula), 633, 635	1000
Paralithodes camtschatica. see red king crab	degradation, impacts on pipelines, 932–933, 936
Paralithodes camtschaticus. see king crab	degradation, impacts on runoff, 917
paralytic shellfish poisoning, 879, 880	degradation, impacts on sanitation infrastructure, 888
Paranoplocephala arctica, 249	degradation, impacts on transportation, 930–931, 935–936
parasites. see also cryptosporidium; echinococcus; Giardia;	degradation, impacts on water supply, 885–886
trichinosis	degradation and boreal forests, 794, 804, 823, 828
and emigration of new species, 386, 419, 436	degradation and carbon storage, 319
of freshwater fish, 412–413	degradation and coastal erosion, 921, 940, 999
impacts of UV radiation, 413	degradation and contaminant release, 430, 432–433, 891
projected increase in aquatic birds, 419, 436	degradation and ecosystems, 550
projected increase in aquatic mammals, 419, 436	degradation and freshwater ecosystems, 998
projected increase in fish, 415, 436	degradation and hazardous materials, 927
parasitic jaeger (Stercorarius parasiticus), 361, 630	degradation and human health, 897
Parasyrphus tarsatus, 258	degradation and landslides, 917, 919–920
Parus caeruleus. see blue tit	degradation and methane emissions, 998
peas, green, 807, 810, 811, 852	degradation and soil subsidence, 916
peas, seed, 810, 811	degradation and subsistence activities, 883-884
peatlands. see also freshwater ecosystems; wetlands	degradation and thermohaline circulation, 999–1000
area of, 363, 562	degradation and transportation, 813
carbon storage, 841, 1014	discontinuous, defined, 209
drainage impacts on biodiversity, 562	and energy balance, 318
in Finland, 795	engineering concerns, 914–916
impacts of projected change in freshwater runoff, 229	feedbacks to global climate, 209, 994
impacts of projected change in permafrost, 215	and groundwater, 45, 357–58
impacts of projected change in river and lake ice, 223	and human activities, 324
prehistoric extent, 252	impacts of degradation, 210
projected change in distribution, 392	impacts of projected change, 219, 588, 677–678, 940
projected impacts, 32, 254, 435, 843	impacts of thaw on animals, 279
water balance, 32	impacts of thawing on indigenous peoples, 667–670
and wildfires, 843	impacts of wildfires, 835
Pechora Basin, 187–188, 227, 229	indigenous observations of change, 279, 993
pectoral sandpiper (Calidris melanotos), 288	and infrastructure, 909–917
Pedicularis hirsuta. see hairy lousewort	methane emissions, 999
Pedicularis spp., 293	observed change, 45, 210–211, 659–660, 676, 910, 913, 963,
pelagic cormorant (Phalacrocorax pelagicus), 626, 635	991, 1005, 1009
pelagic ecosystem	prehistoric extent, 252
defined, 479	present-day distribution, 209, 245
Oscillating Control Hypothesis, 495	projected circumpolar impacts, 211–214
projected impacts, 544	projected impacts, 3, 45, 254, 327, 520, 570, 659–660, 676,
Perca flavescens. see yellow perch	912–913, 952, 990, <i>991</i> , 996–997, <i>997</i> , 1008–1009
Perca fluviatilis. see European perch perch, 71. see also individual species	1012
1	projected regional impacts, 214–215
Percopsis omiscomaycus. see trout perch peregrine falcon (Falco peregrinus), 258, 566	research recommendations, 215–216 and river chemistry, 45
peripheral circulatory diseases, 873	and river chemistry, 45 and river flows, 45
permafrost, coastal and subsea	role of vegetation, 550
background, 216–217	sensitivity to change, 245
and coastal erosion, 519	and snow, 198
impacts of projected change, 215	sporadic, defined, 209
observed change, 217–218	as water source, 547
and offshore engineering, 215	persistent organic pollutants, 878, 947, 952, 982. see also
projected impacts, 188, 218–219, 1008	contaminants
research recommendations, 219–220	analyses, 959

conventions on, 14–15	pipelines. see oil and gas development
and human health, 890–892, 953, 964	place names, 66, 84, 655–656
impacts of climate change on, 427–432	plaice (Pleuronectes platessa), 712
impacts on biodiversity, 566	plankton. see also phytoplankton
in reindeer meat, 980	projected range shifts, 588
Stockholm Convention, 64	plants. see also boreal forest ecosystems; terrestrial ecosystems;
Phaeocystis pouchetii, 481, 482, 504, 506. see also phytoplankton	vegetation
Phalacrocorax aristotelis. see European shag	adaptation to UV radiation, 998
Phalacrocorax carbo. see great cormorant	adaptations to arctic environment, 263–266
Phalacrocorax pelagicus. see pelagic cormorant	change in phenology, 571
Phalaropus fulicaria. see red phalarope	community response to experimental manipulation, 295–296
Phalaropus lobatus. see northern phalarope	diversity, present day, 255–259, 1016
phenols, 624	diversity, projected change, 257
Philomachus pugnax. see ruff	diversity and temperature, 256–259
Phippsia algida. see ice grass	documentation of change, 585
Phleum alpinum. see alpine timothy	genetic diversity, 582
Phoca fasciata. see ribbon seal	genetic response to change, 285–286
Phoca hispida. see ringed seal	impacts of CO <sub>2</sub> concentration change, 275–276
Phoca largha. see spotted seal	impacts of projected change on diseases, 302
Phoca vitulina. see harbour seal	impacts of projected change on human health, 879–892
Phocoena phocoena. see harbour porpoise	impacts of recent climate change, 271–272
Phocoenoides dalli. see Dall's porpoise	impacts of reindeer grazing, 601
Phoebastria nigripes. see black-footed albatross	impacts of UV radiation, 248, 264, 276–277, 953
phosphorus (P)	inventory needs, 580
pollution and biodiversity, 564–565	knowledge of arctic species, 540
in terrestrial ecosystems, 305–309	observed change in distribution, 287
Phyllocnistis populiella. see aspen leaf miner	projected change in distribution, 288–290
Phylloscopus borealis. see Arctic warbler	projected impacts, 272–279, 274–275, 277–278, 998
Phylloscopus inornatus. see yellow-browed warbler	species numbers, 580
phylogeography, 249	tissue chemistry and herbivory, 298–299
Physeter catodon. see sperm whale	platinum. see mineral extraction
phytophagous insects, 563	Plectrophenax nivalis. see snow bunting
phytoplankton	Pleurogrammus monoptergius. see Atka mackerel
bloom onset, 480–481, 491–492, 495, 544	Pleuronectes ferrugineus. see flounder
and carbon cycle, 518	Pleuronectes glacialis. see Arctic flounder
impacts of past climate variability, 497	Pleuronectes platessa, see plaice
impacts of UV radiation, 513, 514	Pleuronectes quadrituberculatus. see Alaska plaice
marine biomass, 479	Pleurozium shreberi, 258
in marine ecosystems, 481	Pluvialis dominica. see lesser golden plover
marine productivity, 479	Pluvialis fulva. see Pacific golden plover
projected change in distribution, 569	Pluvialis squatarola. see black-bellied plover
projected impacts, 504, 505–506, 546, 570, 758	Poa abbreviata. see northern bluegrass
role in marine food webs, 544	Poaceae, 293
Picea abies. see Norway spruce	Pogonatum alpinum, 247
Picea glauca. see white spruce	polar amplification, 23, 38–39, 54, 606, 658
Picea mariana. see black spruce	polar bear (Ursus maritimus), 487, 556, 570
Picea obovata. see Siberian spruce	in Bering Sea, 752
Picea sitchensis. see Sitka spruce	biodiversity monitoring, 586
pigeon guillemot (Cepphus columba), 576, 635	biomass, 479, 480
pilot whale (Globicephala melaena)	harvest by indigenous peoples, 652, 667–668, 668, 674–675
harvest by indigenous peoples, 652	harvest management in Canada, 612, 626
harvest of, 622, 629	historic harvesting, 699
in northwest Atlantic, 732	and human health, 882
population, 736	impacts of past change in sea ice, 674
pine ( <i>Pinus</i> spp.), 262, 551, 784. see also individual species	impacts of POPs, 566
diprionid sawfly outbreaks, 835	importance of sea ice, 496–497, 632
impacts of UV radiation, 852–853	importance to tourism, 600
indigenous observations of change, 80	indigenous observations of change, 70–71, 585, 660, 667
past distribution, 368	and ivory gull distribution, 512
projected impacts, 1006	as keystone predator, 545
in Russian boreal forest, 790	management agreements, 623, 625, 626, 628, 632
treeline change, 1005	migration, 546
Pinguicula spp., 549	multiple stresses on, 615
Pinguinus impennis. see great auk	observed change, 504
pink-footed goose (Anser brachyrhynchus), 288, 574, 1005	overharvesting, 487, 560–561, 1016
pink salmon (Oncorhynchus gorbuscha), 410, 752, 760	paleoecology, 251
pintail, 258	population in Canada, 612
pintail (Anas acuta), 287, 669	productivity, 479
Pinus contorta. see lodgepole pine	projected impacts, 195, 380, 504, 509, 520, 545, 570, 659, 675
Pinus pumila. see dwarf Siberian pine	<i>999, 1004,</i> 1005, 1013, 1017
Pinus sibirica. see Siberian stone pine	and seal prey, 496
Pinus sylvestris, see Scots pine	sport hunting of, 657

tracking, 631	Populus tremuloides. see aspen
polar cod (Boreogadus saida), 485–486	Populus trichocarpa. see black cottonwood
in Bering Sea, 748, 749	Porcupine Caribou Management Board, 637
bycatch, 734	porpoises, 710. see also Dall's porpoise; harbour porpoise
commercial fishery, 652, 693, 697, 748, 761	Potamogeton spp. see pondweed
diet, 480	potatoes, 807, 809, 810, 811, 813
distribution, 484, 486	powan. see European whitefish
harvest by indigenous peoples, 652, 667	precipitation. see also freezing rain; rain-on-snow events; snow
impacts of past climate variability, 738, 739	and avalanches, 917
as keystone predator, 545	in boreal forests, 787–788, 794
in Newfoundland and Labrador, 731	impacts of projected change in sea ice, 195
in northeast Atlantic, 696	impacts of projected change on avalanches, 918–919
in northwest Atlantic, 732	impacts of projected change on breakup timing, 222
as prey, 480, 488, 489–490, 712	impacts of projected change on carbon cycle, 518
projected change in distribution, 545	impacts of projected change on caribou, 615, 661, 668, 670,
projected impacts, 385, 504, 507–508, 509, 740, 759	675
polar desert ecosystems	impacts of projected change on contaminants, 428–430, 545,
biodiversity conservation, 548–551	891
characteristics of, 244-247, 292-293	impacts of projected change on flooding, 917
defined, 11	impacts of projected change on freshwater ecosystems, 547
distribution in Russia, 790	impacts of projected change on human health, 869, 881
net primary production, 306	impacts of projected change on hunting, 617
plant biomass, 306	impacts of projected change on infrastructure, 889, 927
projected change in area, 295, 321	impacts of projected change on landslides, 917–918, 919–920
projected impacts, 326, 998, 1004, 1012	impacts of projected change on muskox, 615, 668
soil characteristics, 255	impacts of projected change on plants, 274–275
soil organic matter, 306	impacts of projected change on polar bears, 545
and temperature, 287	impacts of projected change on river discharge, 923
and treeline advance, 550	impacts of projected change on seabirds, 512
polar fox. see Arctic fox	impacts of projected change on subsistence activities, 667
polar stratospheric clouds, 156, 995	impacts of projected change on thermohaline circulation,
difficulty in simulating, 173	999–1000
impacts of climate change, 419	impacts of projected change on water supply, 885–886
and ozone depletion, 143, 155, 156–157, 419	impacts of projected change on wildfires, 840
and stratospheric temperature, 155, 157, 170–171	indigenous observations of change, 660, 993
and water vapor, 155	influence on tree rings, 819–820
Polar Ural mountains, 849–850	measurement limitations, 31, 39, 185
polar vortex, 156	observational data for model evaluation, 112–113, 114,
and atmospheric circulation, 157	115–116
impacts of climate change on, 419	observed trends, 23, 39-42, 42, 248, 274, 972, 991-992, 1011
and ozone depletion, 154, 169–171, 419	observed variability, 972–973
and ozone levels, 159	present-day, 10, 245
polar willow (Salix polaris), 257, 258, 294–295	projected change, 3, 119, 125, 126–127, 128–129, 133–134,
Pollachius virens. see saithe	139–140, <i>321</i> , <i>326</i> , 470, 661, 812, 917, 918, 952,
pollock. see walleye pollock	972, 994, 1005, 1008, 1011–1012
pollution. see contaminants	and wildfires, 835
polychaetes, 483, 487, <i>504</i>	precipitation and evaporation
polychlorinated biphenyls (PCBs), 877–878, 952, 953, 964. see	projected change, 133–134
also persistent organic pollutants	precipitation and evapotranspiration
analyses, 959	background, 184–186
and arctic fish, 381	estimates of, 185–186
impacts on biodiversity, 565	impacts of projected change, 188
polycyclic aromatic hydrocarbons (PAHs), 381, 877–878, 953	observed trends, 186–187
polygons, 245, 255	projected change, 187–188, 1008
Polygonum spp., 262	research recommendations, 188–189
Polygonum viviparum. see alpine bistort	predator/prey match/mismatch, 494, 496, 505, 505, 506–507, 520
polynyas, 457, 1016	544, <i>570</i> , <i>571</i>
importance to marine mammals, 623–624	and carbon cycle, 518
importance to marine mariniars, 023–024	uncertainties, 521
*	Pribilof Islands
importance to whales, 489 indigenous observations of change, 662	
6	indigenous knowledge case study, 76–77
and marine ecosystems, 483	seabird breeding grounds, 489
productivity, 491	primary production, 544
projected impacts, 477	primnesiophytes, 481
Polysticta stelleri. see Steller's eider	Pristiphora erichsonii. see larch sawfly
Polytrichum commune, 285	productivity
Polytrichum juniperinum, 258	of freshwater ecosystems, 11–12
pond smelt (Hypomesus olidus ), 395	impacts of UV radiation, 425–427
pondweed (Potamogeton spp.), 363	limits on, 543
poplar ( <i>Populus</i> spp.), 784. see also individual species	of marine ecosystems, 479–480
Populus balsamifera. see balsam poplar	projected impacts, 418, 541
Populus tremula. see European aspen	of terrestrial ecosystems, 246

Prosopium cylindraceum. see round whitefish protected areas, 621, 1017–1018	rain-on-snow events. see also precipitation impacts of, 43
in Alaska, 793	observed trends, 42–44
area of, 603	projected increase in, 281
and biodiversity conservation, 541, 573–574, 579, 589–590 in Canada, 604, 791–792	rainbow smelt. see smelt
	rainbow trout (Oncorhynchus mykiss), 407, 702–703, 716, 1012
criteria for selecting, 541–542, 603–604 design approaches, 577–579, 999	Ramsar Convention on Wetlands, 564  Rana spp. see frogs
existing, 570	Rana sylvatica. see wood frog
in Finland, 795	Rana temporaria. see common frog
in Iceland, 800	Rangifer tarandus. see caribou; reindeer; reindeer/caribou
impacts of projected climate change on management, 577–579	Ranunculus glacialis. see glacier buttercup
locations, 603	Ranunculus sabinei. see Sardinian buttercup
management approaches, 578–579	rapeseed. see canola
in Norway, 798	raptors ( <i>Buteo</i> spp.), 268
projected impacts, 570, 1001	Rattus norvegicus. see Norway rat
in Russia, 606–607, 609, 624, 791	raven (Corvus corax), 258, 259, 267
in Sweden, 796	razorbill (Alca torda), 627, 629, 630
and vegetation change, 328	reanalyses. see climate data, reanalyses
and wildlife management, 600, 602–605, 638	recreation, 957
protozoans, 481–482, 540	and boreal forests, 794, 796, 797–798
proxy climate records, 817. see also paleoclimate	impacts on biodiversity, 566–567
and climate scenarios, 101	projected impacts, 200, 1004, 1007
interpreting, 46	recruitment
isotope analysis, 366	definition, 495
tree rings, 366, 814–820, 822–824	red-breasted goose (Branta ruficollis), 288, 289, 574
Psetta maxima. see turbot	red-breasted merganser (Mergus serrator), 629-630, 630
Pseudomonas, 260, 283	red deer (Cervus elaphus), 573
Psora decipiens, 258	red-faced cormorant, 635
ptarmigan (Lagopus mutus), 258, 281	red fox (Vulpes vulpes), 259, 281, 619
adaptation to arctic environment, 267–268	displacement of Arctic fox, 548
harvest by indigenous peoples, 667	impacts of late freeze-up, 76
harvest in Fennoscandia, 617	observed change, 1005
harvest in Russia, 607	red gorgonian (Paragorgia arborea), 561
indigenous observations of change, 70–71, 86, 89	red king crab (Paralithodes camtschatica), 483
and predation, 302	commercial fishery, 749, 763, 767–768
pteropods, 482	impacts of past climate variability, 499, 503, 756–757
Pterostichus costatus, 258	management regimes, 765
Ptilidium ciliare, 258	predation by salmon, 747
Puccinellia phryganodes. see creeping alkaligrass	projected impacts, 759
puffin (Fratercula spp.), 512, 626, 629. see also individual species	red knot (Calidris canutus), 250–251, 258, 288, 290, 361
Puffinus gravis. see greater shearwater	red-legged kittiwake (Rissa brevirostris), 489, 633, 635,
Puffinus griseus. see sooty shearwater	1009–1010
Puffinus tenuirostris. see short-tailed shearwater	red-necked phalarope. see northern phalarope
pulp mills, 565 Puma concolor: see cougar	red-necked stint (Calidris ruficollis), 288, 361 red phalarope (Phalaropus fulicaria), 361, 489, 630
Pungitius pungitius. see ninespine stickleback	red squirrel (Tamiasciurus hudsonicus), 286
Purnumukka, Finland	red-throated loon (Gavia stellata), 363, 633, 635
indigenous knowledge case study, 84–87	
purple sandpiper (Calidris maritima), 630	"redfeed" copepod, 696 redfish ( <i>Sebastes</i> spp.), 486, 767
purple saniapiper ( <i>Cantaris martuma</i> ), 030 purple saxifrage, 258, 272	in Central North Atlantic, 711
purple saxifrage (Saxifraga oppositifolia), 285, 582	commercial fishery, 698, 714, 715
pygmy saxifrage (Saxifraga hyperborea), 258	distribution, 484
Pylaiella littoralis, 480	historic fisheries, 712
1 yidiena ilitoralis, 400	in northeast Atlantic, 696
Q	as prey, 488
Qaanaaq, Greenland	redwing (Turdus iliacus), 258, 260
indigenous knowledge case study, 84	Region 1
qaaqtaq. see least cisco	defined, 1003
Qamani'tuaq. see Baker Lake	fish, number of freshwater and diadromous species, 396–397
Qikiktagrugmiut	freshwater fish populations, projected change in, 406
impacts of late freeze-up, 74–75	Holocene climate and freshwater ecosystems, 367–368
indigenous knowledge case study, 73–76	projected change, 327–329, 1004–1007
qungalik. see Arctic sorrel	projected vegetation change, 326
18	Region 2
R	defined, 1003
rabbit	fish, number of freshwater and diadromous species, 396–397
harvest by indigenous peoples, 654	freshwater fish populations, projected change in, 406–407
and tularemia, 881	Holocene climate and freshwater ecosystems, 368–369
rabies, 880, 881	projected change, 327–329, 1004, 1007–1009
Racomitrium spp., 258, 294	projected vegetation change, 326
radionuclides 565	<del>-</del>

Region 3	and UV radiation, 268
defined, 1003	reindeer husbandry, 660–661, 794–795, 796, 807–808, 813, 1001
fish, number of freshwater and diadromous species, 396–397	coping capacity, 974–976
freshwater fish populations, projected change in, 407	coping constraints, 976–980
Holocene climate and freshwater ecosystems, 369	
	global statistics, 965–968
projected change, 327–329, 1004, 1009–1011	and government policy, 978
projected vegetation change, 326	and habitat loss, 976–977
Region 4	impacts of industrial development, 608
defined, 1003	impacts of treeline change, 651
fish, number of freshwater and diadromous species, 396–397	impacts of vegetation change, 659
freshwater fish populations, projected change in, 407–408	impacts on treeline, 851
Holocene climate and freshwater ecosystems, 370–371	impacts on wildlife, 601
projected change, 327–329, 1004, 1011–1014	by indigenous peoples of the Russian North, 678–680
projected vegetation change, 326	and legal structures, 978–979
regional climate models (RCMs), 103. see also downscaling	multiple stresses on, 970–971, 980
	1 /
of arctic atmosphere, 130–134	and overgrazing, 609, 657–658, 676–678, 680, 956
of Arctic Ocean, 135	and pollution, 979–980
coupled arctic models, 135–136	and predator control, 616, 619, 977
HIRHAM, 130–132, 133, 135–136	projected impacts, 558, 948, 969, 1000, 1003, 1004, 1007
for improving projections, 142	in Russia, 609, 664
projections of precipitation change, 133–134	by Saami, 969–970
projections of pressure change, 134–135	and Saami culture, 894
projections of temperature change, 133–134	and snow quality, 958, 973, 975, 1007
simulation of present-day climate, 131–132, 135–136	technological change, 952
uncertainties in projections, 120	vulnerability case study, 948, 965–980
	and wildlife management, 637
regional impacts, 1018–1019. see also Region 1; Region 2; Region	8 /
3; Region 4	by Yamal Nenets, 676–678
reindeer	Reinhardtius hippoglossoides. see Greenland halibut
Svalbard reindeer, 969	research recommendations
tundra reindeer, 969	agriculture, 810, 854
wild forest reindeer, 969	biodiversity, 580–589
· · · · · · · · · · · · · · · · · · ·	
reindeer (Rangifer tarandus), 252. see also reindeer/caribou	boreal forests, 854
commercial hunting in Russia, 608–610	co-management, 686–687
cultural importance, 655	coastal and subsea permafrost, 219–220
distribution, 968	coastal erosion, 921–922
harvest in Russia, 607	commercial fisheries, 771–772
impacts of herding on wild populations, 601	
	fisheries management, 705, 767
impacts of infrastructure development, 567	freshwater discharge, 229–230
impacts of overgrazing, 562–563, 601	freshwater ecosystems, 437–438
impacts of permafrost thaw, 660	future assessments, 1018–1019
impacts of warm winters, 1005	glaciers and ice sheets, 208–209
indigenous observations of change, 85–90, 279	,
	human health, 875, 877, 878, 897, 900–902
management in Russia, 610–611	impacts on indigenous peoples, 686–687
observed change in wild populations, 1005	improving model projections, 140–144
overgrazing, 657	indigenous knowledge, 93–94, 901
population declines in Russia, 609	indigenous peoples adaptation, 665
predators, 977	
1 /	indigenous subsidence activities, 650–652
projected impacts, 611, 676, 973–974	infrastructure, 915, 941
reindeer/caribou (Rangifer tarandus), 258, 556, 1016. see also	marine Arctic, 520–522
caribou; reindeer	marine mammals, 631
adaptation to arctic environment, 266–267, 556–557	ozone and UV radiation, 175–176
adaptations to arctic climate, 966–967	permafrost, 215–216
	1 /
biodiversity monitoring, 586	precipitation and evapotranspiration, 188-189
distribution, 969	river and lake ice, 223, 923
harvest by indigenous peoples, 652–653	sea ice, 195–196
hunting of, 247	sea-level rise, 236
impacts of ice crusting, 279, 280	seabird management, 628, 633–634, 636
impacts of projected change in snow, 301, 660	snow, 200–201
impacts on vegetation, 300	societal conditions and trends, 983
importance to indigenous peoples, 599–600	vegetation change and reindeer husbandry, 678
indigenous observations of change, 280	vulnerability assessments, 982–983
and lichens, 273	wildlife management, 621, 638, 639
	2 , , ,
management, 637–638, 639	resilience
migration triggers, 267	of arctic communities, 658, 709
monitoring, 618	assessing, 9
observed change, 661	definition, 950
overharvesting, 560	of human-environment systems, 946–949, 951
parasites and climate change, 302–303	and human health, 865
population cycles, 268	of indigenous peoples, 92–93, 670, 686
and predation, 301–302	and multiple stresses, 15
projected impacts, 281, 604, 659, 660–661, 998	research recommendations, 902
snow beds as refuge, 275	and wildlife co-management, 666

resource development. see also forestry; hydroelectric generation;	sensitivity to temperature change, 32
mineral extraction; oil and gas development	types of, 31–32
impacts of projected change in freshwater runoff, 229	and wood transport, 786
impacts of projected change in permafrost, 215	roach (Rutilus rutilus), 406, 513-514
impacts on arctic ecosystems, 605	rock sole (Lepidopsetta bilineata), 487
impacts on indigenous peoples, 651	commercial fishery, 751
and improved transportation, 923 respiratory diseases, 869, 872–873	impacts of past climate variability, 502, 503, 503 projected impacts, 508
Rhizoplaca melanophthalma, 269	rockfish. see redfish (Sebastes spp.)
Rhodostethia rosea. see Ross' gull	Ross' gull (Rhodostethia rosea), 288, 490
rhubarb, wild	rotatorians, 480
harvest by indigenous peoples, 654	rough-legged buzzard (Buteo lagopus), 301
Rhytidiadelphus triquetrus, 258	round whitefish (Prosopium cylindraceum), 378
ribbon seal ( <i>Phoca fasciata</i> ), 488–489	rowan (Sorbus aucuparia), 563, 799
commercial harvest, 753 harvest of, <i>625</i>	Rubus chamaemorus. see cloudberry ruddy turnstone (Arenaria interpres), 250–251, 361, 630
projected impacts, 504, 510	ruff (Philomachus pugnax), 287
right whale (Eubalaena glacialis), 489	ruffe (Gymnocephalus cernuus), 406, 576
ring-necked duck (Aythya collaris), 363	Russia
ringed seal (Phoca hispida), 488, 624	commercial fisheries, 695–709
in Bering Sea, 752	demographics, 868–869
biodiversity monitoring, 586 commercial harvest, 715, 753	economic importance of fisheries, 700–704 fisheries management, 704–706
cultural importance to Inuit, 674	forest area, 784
harvest by indigenous peoples, 652, 667–668, 671, 672, 674,	forest tenure and management, 789–791
882	sanitation systems, 884
harvest of, 622, 625, 627, 628, 632	wildfire regimes and management, 836-837
impacts of past change in sea ice, 674	wildlife management, 606–610, 623–626, 638
indigenous observations of change, 71, 670	Russian Association of Indigenous Peoples of the North
migration, 546 and persistent organic pollutants, 953	(RAIPON), 13, 68–71, 87, 602 Russian North, 678–680, 968
as prey, 487, 615	Russian Sea Route. see Northern Sea Route
projected impacts, 504, 509–510, 520, 545, 570, 675, 964,	Rutilus rutilus, see roach
1005, 1011	
Rissa brevirostris. see red-legged kittiwake	S
Rissa tridactyla. see black-legged kittiwake	Saami, 599, 616, 652
river discharge, 223–230. see also rivers	historic adaptations, 662
and contaminant transport, 891, 947 estimated, 225	impacts of hydroelectric development, 657 indigenous knowledge, 960
and hydroelectric generation, 188	observations of change, 84–90, 271–272, 279, 660, 679
impacts of permafrost change, 913	projected impacts, 1007
impacts on thermohaline circulation, 999-1000	reindeer herding, 658, 661, 664, 794–795, 894, 948, 956,
and navigation, 188, 1000	965–968, 981
observed change, 997	and technological change, 957
projected change, 127, 137–138, 188, 917, 923, 997, 1004, 1008, 1012	terminology for snow, 958, 973 vulnerability case study, 962
and projected change in river and lake ice, 223	Saami Council, 13, 68–71, 602, 968
and projected change in snow cover, 199–200	Saami knowledge. see indigenous knowledge
and water supply, 1000	Sabine's gull (Xema sabini), 489, 630
river otter, 619	sable (Martes zibellina), 607
rivers. see also freshwater ecosystems; ice, river and lake; specific	sablefish, 763, 765, 769
rivers breakup and freshwater ecosystems, 998	Sachs Harbour, NWT, Canada
characteristics of Siberian, 546–547	impacts case study, 666–671 indigenous observations of change, 69, 71, 666–671
food webs in, 359	vulnerability case study, 961–963
freshwater ecosystems of, 358–360	saffron cod (Eleginus gracilis), 74, 75, 760
impacts of changing UV radiation, 424-425	Sahtú, 81
impacts of freeze-up timing change, 611	saithe (Pollachius virens)
impacts of observed temperature change, 23, 32	export value, 704
impacts of permafrost thaw, 45 impacts of projected change on anadromous fish, 398–399	impacts of past climate variability, 717 in Northeast Atlantic, 696
impacts of projected change on anadromous hish, 398–399	past distribution change, 500
indigenous observations of change, 88–90, 685	as prey, 696
observed change in ice cover duration, 997	salinity
projected change in flow rates and timing, 376-377, 379	impacts of projected change in freshwater runoff, 228, 99
projected change in freeze-up timing, 374	impacts of projected change in sea ice, 195
projected change in ice cover duration, 659, 997	impacts of projected change on marine ecosystems, 659 impacts of projected sea level rise, 235
projected change in thermal gradient, 375 projected change in timing and severity of breakup, 375, 380,	observed change, 44–45, 467–469, 720
434	projected impacts, 476, 478
projected change in timing of spring freshet, 374–375	Salix arctica. see Arctic willow
projected impacts 379–381 434 436	Salir planca see plancous willow

Salix herbacea. see dwarf willow	projected change in distribution, 584
Salix lanata. see hairy willow	projected impacts, 824
Salix myrsinifolia. see dark-leaved willow	in Sweden, 797
Salix myrsinites. see whortle-leaved willow	tree ring data, 819–820
Salix myrtilloides, 258	sculpin (Cottus spp.). see also fourhorn sculpin; slimy sculpin
Salix phylicifolia. see tea-leaved willow	bycatch, 761
Salix polaris. see polar willow	commercial fishery, 761
Salix pulchra. see tealeaf willow	impacts of past climate variability, 756
Salix reptans, 258	as prey, 488
Salix spp., 292	projected impacts, 758, 759
Salmo salar. see Atlantic salmon	sea bass (Dicentrachus labrax), 716
Salmo trutta. see brown trout	sea ducks, 483
salmon (Oncorhynchus spp., Salmo salar). see also individual	sea ice. see also land-fast ice
species	and accessibility to resources, 565-566, 883, 964, 965
aguaculture, 695, 708, 716, 747	albedo feedback, 474, 994
and botulism, 880	albedo measurements, 190
commercial harvest, 657	area, seasonal change in, 30
	The state of the s
export value, 704	background, 189–190
harvest by indigenous peoples, 652, 654	in Bering Sea, 755
impacts of past climate variability, 503, 658	and Bering Sea fish, 502–503
indigenous observations of change, 585, 880	and carbon exchange, 515–517
observed change, 1009–1010	characteristics, 456–458
and permafrost degradation, 884	climate model representation, 107–108
projected impacts, 507, 660, 1010	and coastal erosion, 76–77, 219, 234
salmon shark, 77	and contaminants, 954
	*
salmonids. see also fish, anadromous; salmon; individual species	coverage, 189–190, 454
diversity of, 549	distribution and thickness, 189–190, 457
and fisheries, 412	and DMS, 143
impacts of past climate variability, 503	drift patterns, 44, 457–458
impacts on habitat, 402	ecosystems, 480, 482, 491, 506, 659
potential distribution change, 410	extent, observed trends, 35, 37
species, number of arctic, 396–397	first-year distribution, 456
salt marshes. see wetlands	freeze-up timing, impacts of, 74–75
	• • • · · · · · · · · · · · · · · · · ·
Salvelinus alpinus. see Arctic char	and freshwater influx, 208
Salvelinus confluentus. see bull trout	impacts of precipitation change, 188
Salvelinus fontinalis. see brook trout	impacts of projected change, 194–195, 195
Salvelinus malma. see Dolly Varden	impacts of projected change in freshwater runoff, 228
Salvelinus namaycush. see lake trout	impacts of projected change in snow cover, 200
Sami. see Saami	impacts of projected change on aquaculture, 745
sand eel. see sand lance	impacts of projected change on fisheries, 741
sand lance ( <i>Ammodytes</i> spp.), 488, 712, 732	impacts of projected change on human health, 882–883
	impacts of projected change on indigenous peoples, 659,
Sander lucioperca. see zander	
Sander vitreus. see walleye	667–670, 670
sanderling (Calidris alba), 258, 259, 288, 289	impacts of projected change on infrastructure, 888-889
sandhill crane, 70	impacts of projected change on marine ecosystems, 520
sandpipers (Calidris spp.), 259, 268, 358. see also individual	impacts of projected change on nutrient supply, 544
species	impacts of projected change on seals, 740–741
Sanguisorba officinalis. see official burnet	impacts of temperature change, 3
Sapmi	impacts on marine ecosystems, 505
indigenous knowledge case study, 84–87	impacts on reindeer, 609
sardine, 508	importance to marine ecosystems, 490, 623–624
Sardinian buttercup (Ranunculus sabinei), 259	importance to polar bears, 487, 504
sawflies, 259, 268, 549	importance to seals, 488
Saxifraga caespitosa. see tufted alpine saxifrage	importance to walrus, 488
Saxifraga cernua. see nodding saxifrage	indigenous observations of change, 68-69, 84, 94, 190, 632
Saxifraga hyperborea. see pygmy saxifrage	660, 661–662, 882, 886, 993
Saxifraga nivalis. see alpine saxifrage	during Little Ice Age, 250
Saxifraga oppositifolia. see purple saxifrage	melt season duration, 457
	· · · · · · · · · · · · · · · · · · ·
Saxifragaceae, 293	multi-year distribution, 456
scallops, 561. see also Iceland scallop	observational data for model evaluation, 118
Scandinavia	observed change, 190–192, 467, 659, 963
observed change in glaciers and ice sheets, 206	and ocean mixing, 460–461
scenario. see climate scenarios; emissions scenario	perennial, defined, 30
Scoloplos armiger, 483	perennial, observed trends, 37, 190–192
Scomber japonicus. see Pacific mackerel	projected impacts, 3, 122, 192–194, 480, 519, 659, 964
Scomberesox saurus. see Atlantic saury	research recommendations, 195–196
scoters ( <i>Melanitta</i> spp.), 418, 627. see also individual species	seasonal, defined, 30
Scots pine (Pinus sylvestris), 278	seasonal, projected change, 472
climate responses, 822–824, 828–830	seasonal cycle, 456–457
in Finland, 795	and species exploitation, 560
impacts of UV radiation, 852–853	and subsistence activities, 660
observed change, 829–830	thickness, observed trends, 30, 37, 191–192, 457

thickness, projected change, 472	impacts on water supply, 885
trends in Russian Arctic, 624	impacts on water treatment, 885
and UV radiation, 174	isostatic, 230
sea ice duration	observed, 232–233, 997
and ENSO, 469	and polar ice caps, 34
and fish migration, 496	projected, 33–34, 233–235, 570, 994, 997
impacts of observed change on polar bear health, 615	projected contribution of glaciers and ice sheets, 207–208
impacts of projected change, 472–475	research recommendations, 236
impacts of projected change on marine mammals, 496–497	steric, 230–231
impacts of projected change on polar bears, 545, 674	sea otter (Enhydra lutris), 489
impacts of projected change on ringed seals, 674	in Bering Sea, 752
and NAO, 469	historic harvesting, 761
and PDO, 469	management agreements, 632
and phytoplankton bloom, 491–492, 495	overharvesting, 487, 632–633, 1016
projected impacts, 476, 570	projected impacts, 510, 759
sea ice extent	sea-tangle (Fucus distichus), 480, 481
and the Arctic Oscillation, 44	seabirds, 487–490. <i>see also</i> birds; birds, aquatic; birds, migratory;
and contaminant transport, 891, 947	waders; individual species
impacts of projected change on coastal erosion, 922, 940, 952,	in Bering Sea, 746
999	biodiversity monitoring, 586
impacts of projected change on marine ecosystems, 544, 569,	biomass, 479, 480
588, 659, 694, 952	breeding areas and sea ice extent, 497
impacts of projected change on marine mammals, 509–511	climate impacts on, 496–497
impacts of projected change on navigation, 922–923	commercial harvest, 715
impacts of projected change on polar bears, 545	diversity, 489
impacts of projected change on productivity, 505, 505-506	food requirement, 484
impacts of projected change on seabirds, 635–636	harvest by indigenous peoples, 710
impacts of projected change on seals, 545	harvest of, 622–623, 625–626, 627–628, 629–631, 633–635,
impacts of projected change on subsistence activities, 656, 922	710
impacts of projected change on transportation, 934–935, 1002	historic ranges, 624
impacts of projected change on travel, 952–953	impacts of Arctic fox, 576
impacts of projected change on water supply, 886	impacts of capelin overharvesting, 624
and northern shrimp, 738–739	impacts of introduced predators, 563, 605
in northwest Atlantic, 731	impacts of oil and gas development, 624
	· · · · · · · · · · · · · · · · · · ·
observations of, 35	impacts of oil pollution, 565
observed change, 23, 44, 54, 190–192, 457, 471, 923–924,	impacts of past climate variability, 504
991, 997, 1009, 1018	importance to indigenous peoples, 622, 634–636
paleoclimatic, 48	importance to tourism, 600
projected impacts, 119, 127, 139, 471–472, 476, 659, 694,	Important Bird Areas, 634
924–926, 952, 990, 991, 996–997, 999, 1005, 1012	indigenous observations of change, 993
Sea-Ice Model Intercomparison Project, 107, 109	indirect impacts of UV radiation, 953
sea ice thickness	influenza, 880
impacts of projected change on infrastructure, 927	management and conservation, 622-636, 765
observed trends, 44	mercury levels in, 566, 964
and subsea permafrost development, 217–218	observed change, 1009
sea level	and ocean temperature, 571–572
early Holocene, 367	overharvesting, 487, 561, 1016
Last Glacial Maximum, 249	productivity, 479, 484
observed change, 33	projected impacts, 504, 509, 511–512, 545–546, 572, 604, 636,
paleoclimatic, 48–49, 253	768, 999, 1005, 1010
projected change, 253	projected range shifts, 588
sea level pressure. see atmospheric pressure	reproductive timing and food supply, 496
sea level rise	research recommendations, 633–634, 636
and atmosphere-ocean processes, 231–232	status, 630, 635
1 1	
attribution, 236	seals, 488–489. see also individual species
background, 230	adaptation to cold, 556
eustatic, 230–231	and animal rights movement, 615, 657, 956, 965
and "freshwater" storage, 34	and benthos, 483
and glaciers and ice sheets, 203, 204–206, 234	biomass, 479
impacts of projected change, 235–236	and botulism, 880
impacts on aquatic ecosystems, 923	commercial harvest, 715
impacts on birds, 290	and contaminants, 964
impacts on coastal ecosystems, 208	food requirement, 484
impacts on coastal erosion, 219, 234, 235–236, 519, 922–923,	harvest, 710
940, 999	harvest by indigenous peoples, 652, 655, 668, 668, 710
impacts on coastal lakes, 383	historic change, 663
impacts on coastal permafrost, 219	historic ranges, 624
impacts on coastal stability, 230–236	and human health, 882
impacts on indigenous peoples, 559	hunting by indigenous peoples, 650
impacts on marine ecosystems, 505, 659	impacts of past climate variability, 504
impacts on protected areas, 589–590	impacts of projected change in sea ice, 659
impacts on protected areas, 507 576 impacts on seabirds, 512, 546, 635–636	importance of sea ice, 496–497, 632
r,,,,	

indigenous observations of change, 70–71, 585, 667	snow. see also precipitation; snow cover
indirect impacts of UV radiation, 953	albedo feedback, 994
and ivory gull distribution, 512	depth and plants, 264
management and indigenous subsistence activities, 665	depth impacts on black spruce, 851
management in Canada, 626–627	depth impacts on plants, 274–275
in northeast Atlantic, 696	depth impacts on ungulates, 570
productivity, 479, 484	duration and bird breeding, 290
projected impacts, 380, 510, 545, 660, 965, 999, 1010, 1013,	duration and plants, 264
1017	duration and subsistence activities, 883
reproduction and food supply timing, 496	duration and UV radiation exposure, 277
sedges, 294. see also individual species	impacts of projected change on biodiversity, 589
sei whale (Balaenoptera borealis), 489, 715, 732, 736	impacts of projected change on caribou, 661
selenium, 890	impacts of projected change on geese, 280
Seligeria polaris, 258	impacts of projected change on indigenous hunting, 683–684
semipalmated plover (Charadrius semipalmatus), 361	impacts of projected change on plants, 274–275
Severnaya Dvina, 226	impacts of projected change on spring freshet, 374–376
Severnaya Dvina Basin, 227	impacts of projected change on subsistence activities, 667, 668
· · · · · · · · · · · · · · · · · · ·	
Severnaya Zemlya	impacts of rain-on-snow events on melt, 43
observed change in glaciers and ice sheets, 206–207	importance to polar bears, 487
sharks, 486. see also individual species	and marine ecosystems, 490
sharp-tailed sandpiper (Calidris acuminata), 288	melt and flooding, 917
shearwater ( <i>Puffinus</i> spp.), 627–628. see also individual species	melt and landslides, 918
Shishmaref, 660, 888–889	observed change, 42
shore lark (Eremophila alpestris), 1006	observed change in depth, 198–199
shorebirds. see waders	projected change, 1008
short-eared owl (Asio flammeus), 301	and reindeer husbandry, 973, 1000
short-finned squid. see squid	and winter roads, 933
short-tailed albatross, 635	snow beds, 290, 300
short-tailed shearwater (Puffinus tenuirostris), 624	importance to reindeer/caribou, 275
shrews (Sorex spp.), 267	indigenous observations of change, 275
shrimp. see also individual species	and vegetation, 274–275
commercial fishery, 742, 761, 964	as water source, 547
harvesting and biodiversity, 561	snow blindness, 152, 168, 996
historic change, 663	snow bunting (Plectrophenax nivalis), 258
historic change, 603 historic harvesting, 762	snow cover
impacts of past climate variability, 766, 866	albedo, 198, 998
predation by Atlantic cod, 739	and avalanches, 918–919
projected change in distribution, 700	background, 196–198
projected impacts, 504, 758, 768	duration and albedo, 322
Siberian fir (Abies sibirica), 252	impacts of projected change, 200
Siberian larch (Larix sibirica), 800, 849	impacts of projected change on caribou/reindeer, 660
climate responses, 822–824	impacts of projected change on subsistence activities, 660
projected impacts, 802, 803	impacts of projected change on underwater UV, 419–420
Siberian silkworm (Dendrolimus sibiricus), 576	impacts of temperature change on duration, 3
Siberian spruce (Picea obovata), 822–824, 848–849	impacts of thaws on duration, 43
Siberian stone pine (Pinus sibirica), 252	impacts on marine ecosystems, 505
Siberian whitefish (Coregonus pidschian), 395, 397	importance to climate projections, 107
sibling vole (Microtus rossiaemeridionalis), 279, 280	importance to polar bears, 496–497
Sika deer (Cervus nippon), 573	importance to seals, 496–497, 545
sila, 655	indigenous observations of change, 660, 993
Silene acaulis. see moss campion	measurement limitations, 196
silver birch (Betula pendula), 571, 795, 852, 853	observed change, 198–199, 274, 991, 996–997
Sisimiut, Greenland, 663	observed change in extent, 23, 31, 54, 1009
Sitka spruce ( <i>Picea sitchensis</i> ), 800, 831–833, 845	and permafrost, 909
skates	and permanost, 909 and persistent organic pollutants, 429–430
impacts of past climate variability, 503, 756	present-day distribution, 196–197
projected impacts, 758, 759, 760	projected change, 918, 972, 996–997
skin cancer, 875–876, 878, 899, 953, 996, 1001	projected change in duration, 3, 659
skin rashes, 63	projected change in extent, 119, 122, 127, 199–200, 201
skuas (Stercorarius spp.), 268, 301, 490	research recommendations, 200–201
slides. see landslides	and subsistence activities, 965
slimy sculpin (Cottus cognatus), 378, 564	and UV radiation, 153, 160, 163–164, 174–175, 996
smallmouth bass (Micropterus dolomieu), 398, 405	and vegetative masking, 197–198
smelt (Osmerus mordax), 361, 397	snow crab (Chionoecetes opilio), 483
in Bering Sea, 748	in central North Atlantic, 711
harvest and late freeze-up, 75	commercial fishery, 735, 749, 763, 765, 767
projected impacts, 760	impacts of past climate variability, 499, 739, 757
as traditional food, 74	in northwest Atlantic, 732
smooth bromegrass, 807	observed change, 1002
snails, 483	projected impacts, 740, 768, 1007, 1013
,	snow gentian (Gentiana nivalis), 557
	( 50 ( 50

snow goose (Anser caerulescens), 363	Sphagnum spp., 277, 293, 317, 427, 571
harvest by indigenous peoples, 667, 669	spiders, 580
impacts on salt marsh sward, 300	Spio filicornis, 483
indigenous observations of change, 70–71, 662	Spiochaetopterus typicus, 483
projected habitat loss, 288	sponges, 564, 580, 756
Snow Models Intercomparison Project (SMIP), 109	spoon-billed sandpiper (Eurynorhynchus pygmaeus), 288, 289
snow sheep, 609	spotted redshank (Tringa erythropus), 258
SnowChange project, 79, 84, 87, 960	spotted seal (Phoca largha), 488
snowshoe hare (Lepus americanus), 268, 280, 552, 605, 610, 619	commercial harvest, 753
snowy owl (Nyctea scandiaca), 258, 267, 301, 550	harvest and late freeze-up, 74–75
observed change, 1005	harvest of, 632
projected change, 1006	indigenous observations of change, 71
societal impacts. see arctic residents; indigenous peoples	projected impacts, 504, 510
sockeye salmon	spring wheat, 584
commercial fishery, 752	springtails. see collembolans
harvest by indigenous peoples, 654	spruce (Picea spp.), 784. see also individual species
impacts of past climate variability, 503	indigenous observations of change, 80
observed range change, 410	monocultures in Norway, 798
projected impacts, 760	past distribution, 369
record of past climate change, 658	projected impacts, 833
soil	in Russian boreal forest, 790
acidification in boreal forests, 795	spruce bark beetle (Dendroctonus rufipennis), 552, 831–833, 879
and agriculture, 809	spruce budworm (Choristoneura fumiferana), 833–834
biodiversity, 554	spruce coneworm (Dioryctria reniculelloides), 833–834
carbon balance and warming, 283, 319, 554, 846	spruce engraver beetle ( <i>Ips typographus</i> ), 834
carbon balance and wildfires, 842, 843	spurdog (Squalus acanthias), 500
carbon in boreal forests, 785	Squalus acanthias. see spurdog
carbon storage, 305, 1014	squid (Illex illecebrosus)
characteristics of, 246	commercial fishery, 761
erosion in Iceland, 799–800	in northwest Atlantic, 732
experimental warming and boreal forest growth, 820–821	as prey, 487, 488, 712
impacts of permafrost change on moisture, 913	projected impacts, 740, 759–760
impacts of projected moisture change on biodiversity, 588	St. George Island
nutrient levels, 305	coastal erosion, 77
organic matter content, 306	St. Lawrence Island, Alaska
and plants, 265	indigenous observations of change, 69
projected change in moisture, 188	St. Lawrence Island polynya, 499, 507
species richness, 540	stamukhi (ice ridges), 30. see also land-fast ice
thermal regimes and mosses, 273	starfish, 756
and trace-gas exchange, 311–314	Steller sea cow ( <i>Hydrodamalis gigas</i> ), 487, 560, 601, 1016
and vegetation response, 255	Steller sea lion, 77, 489
soil bearing strength, 912, 915	in Bering Sea, 752–753
solar activity	historic harvesting, 761
· · · · · · · · · · · · · · · · · · ·	management agreements, 632, 767
impacts on Bering Sea ecosystem, 757–758	observed change, 1009
trends, 160 and UV radiation, 160, 160–161, 250	population trends, 633
solar radiation	projected impacts, 510, 759
early Holocene, 367	and walleye pollock fishery, 750
observational data for model evaluation, 117, 118	Steller's eider ( <i>Polysticta stelleri</i> ), 76, 363, 630, 633, 635
and ozone levels, 154	Stenodus leucichthys. see inconnu
projected change, 127	steppe–tundra, 250, 252, 584, 998, 1016
solar zenith angle, 153, 160, 161, 162	Stercorarius longicaudus. see long-tailed jaeger
solifluction, 916	Stercorarius parasiticus. see parasitic jaeger
Somateria fischeri. see spectacled eider	Stereocaulon paschale, 276
Somateria mollissima. see common eider	Sterna aleutica. see Aleutian tern
Somateria spectabilis. see king eider	Sterna hirundo. see common tern
Somniosus macrocephalus. see Greenland shark	Sterna paradisaea. see Arctic tern
sooty shearwater (Puffinus griseus), 732	Stockholm Convention on Persistent Organic Pollutants, 602
Sorbus aucuparia. see rowan	stone loach (Noemacheilus barbatulus), 396
Spartina, 573	stoneflies, 549
species. see also animals; plants; wildlife	storms. see also extreme events; precipitation
adaptation to current environment, 11, 254, 543, 1017	indigenous observations of change, 660
documenting, 580–583	projected change, 470, 918
projected change in abundance, 571–572	stratification, ocean
projected impacts, 254–255, 1017	impacts of projected change in sea ice, 195
species distribution, 568–570	impacts on sea ice, 520
barriers to change, 588, 997, 1012, 1017–1018	projected impacts, 208
observed change in, 608	uncertainties, 521
projected change, 998, 1004	stratosphere
spectacled eider (Somateria fischeri), 363, 489, 633, 635	representation in climate models, 142–143
sperm whale (Physeter catodon), 715, 732, 736, 753	sturgeon, 70, 396
Spermophilus parryii. see ground squirrel	

succession, ecological	and numan nearth, 870–874
definition, 583	impacts of observed change, 23, 32
suicide, 896, 897	impacts of projected change on aquatic birds, 617
Sukachev larch (Larix sukaczewii), 800	impacts of projected change on biodiversity, 588, 589
sulfur, 565	impacts of projected change on buildings, 930
	impacts of projected change on coastal permafrost, 218–219
sulfur oxides, 312	
sunburn, 996	impacts of projected change on frost heave, 916
increases in, 63, 152, 168	impacts of projected change on heating requirements, 936, 937
and UV radiation, 953	impacts of projected change on hunting, 617
sunflower, 807, 810, 812	impacts of projected change on infrastructure, 928
"super-dominant" species, 11, 260, 549, 1017	impacts of projected change on permafrost, 912–913
Surface Heat Budget of the Arctic Ocean (SHEBA), 131, 185	impacts of projected change on pipelines, 932–933
Sustainability of Arctic Communities Synthesis Model, 684–685	impacts of projected change on plants, 272–273
Svalbard	impacts of projected change on seabirds, 511–512, 635–636
observed change in glaciers and ice sheets, 206	impacts of projected change on species ranges, 588
species richness, 580	impacts of projected change on subsistence activities, 667, 883
Svalbard Airport	
	impacts of projected change on vegetation, 967
projected impacts, 939–940	as indicator, 899
Svalbard ptarmigan (Lagopus mutus hyperboreus), 267	indigenous observations of change, 660, 661, 684, 993
swans, 358	and insect outbreaks, 831
Sweden	latitudinal gradients, 11, 257, 373–376
demographics, 868–869	observational data for model evaluation, 111–114, 115
• •	
forest area, 784	observed change, 3, 23, 35–39, 54, 248, 250, 457, 467, 468,
forest tenure and management, 796–798	468, 606, 659, 676, 716, 908–912, 971, 991–992,
human health, 870	1007, 1009, 1011, 1018
projected impacts on boreal forest, 797	observed change in boreal regions, 786, 788
protected areas, 541	paleoclimatic, 253
*	past change, 1018
swordfish (Xiphias gladius)	
past distribution change, 500	present-day, 10, 245
sympagic organism	projected change, 3, 4, 119, 120, 121, 121–125, 128–129,
defined, 480	133–134, 138–139, 248, <i>253, 321, 326,</i> 373–376,
Synthliboramphus antiquus. see ancient murrelet	470, 661, 700, 739–740, 801–806, 805, 806, 917,
symmetric amprims and quasi-see another marrore	918, 938–940, 952, 964, 971–972, 994, 1004–1005,
T	
T	1007, 1009, 1011
taiga. see also boreal forest ecosystems	rate of change, 253–254
defined, 785	reconstruction with tree rings, 814–820
taiga bean goose (Anser fabalis fabialis), 574	and reindeer husbandry, 894
taliks, 914	and sanitation infrastructure, 888
definition of, 357	and seabird distributions, 512
tamarack (Larix laricina), 834	and treeline change, 850
Tamiasciurus hudsonicus. see red squirrel	trends in Russian Arctic, 624
Tanana River, Alaska, 221–222	and water treatment, 885–886
Tanner crab (Chionoecetes bairdi), 483	and wildfires, 803–804, 835–836, 837, 840–841
commercial fishery, 749	and winter roads, 933–934
impacts of past climate variability, 499, 503, 757	
	temperature, ocean
projected impacts, 758, 759–760	and aquaculture, 745
Taraxacum officinale. see dandelion	and carbon balance, 517
Taymir Peninsula, 246, 256, 259, 260, 293, 609, 610, 1016	and climate oscillations, 572
past climate, 815–817	and cod recruitment, 500
1	
projected climate, 802–803	and fish distribution, 496, 520, 545
tea-leaved willow (Salix phylicifolia), 799, 853	and fish production, 544
tealeaf willow (Salix pulchra), 274	and human health, 896–897
temperature	impacts of projected change in sea ice, 195
impacts on freshwater fish, 547	impacts on aquaculture, 708
impacts on migration, 418	impacts on fish, 494–496, 633, 700
· · · · · · · · · · · · · · · · · · ·	•
increases, impacts on contaminants, 428, 432	impacts on marine ecosystems, 659
and marine ecosystems, 490	impacts on marine mammals, 509, 633
and marine productivity, 491	impacts on nutrient supply, 544
and vegetation change, 324	impacts on species ranges, 588
temperature, air. see also air freezing index; air thawing index;	impacts on zoobenthos, 544
diurnal temperature range; rain-on-snow events; thaw	and marine benthos, 497–498
1 0,	,
frequency	and marine ecosystems, 490–491
attribution, 37–38, 39	observed change, 44–45, 468, 716, 720, 1009
and avalanches, 918–919	and PDO, 469
in boreal forests, 794, 805	and predator-prey relationships, 495
and change in disease occurrence, 880	projected change, 472, 475, 476, 519, 694, 1005, 1006
and change in sea ice, 194–195	and red tides, 879
comparison of arctic and global trends, 35	and seabirds, 571–572
and contaminants, 954	and subsea permafrost, 218
and distributions of plants and animals, 879	variability, 467–469
and eastern spruce budworm, 833	and wildfires, 836
and geometrid moths 299	and zoonlankton hiomass 495 497
ADD VEOLUCIO MOIDS 799	AND ZOODIANKION DIOMASS 497 497

temperature, soil	overharvesting, 631
and infrastructure, 910–912, 914–916	status, 630, 635
and methane emissions, 312–313, 317	thorny sculpin (Icelus spiniger), 496, 756
observed change, 910	threespine stickleback (Gasterosteus aculeatus), 385, 409
projected change, 912	thresholds of change
and vegetation cover, 550	in ecosystems, 247
temperature, stratospheric	paleoclimate, 54
impacts of trace gases, 155	Thule culture, 663, 672, 799–800
and ozone depletion, 154, 156, 157, 169–171, 995	Thunnus thynnus, see bluefin tuna
and polar stratospheric clouds, 155	Thyasira sp., 483
projected change, 157, 171, 419, 996	Thymallus arcticus. see Arctic grayling
temperature, water	Thysanoessa spp., 482
and metal accumulation in fish, 429	timothy, 807, 810
terns. see also individual species	tin. see mineral extraction
harvest of, 627–628, 629, 633	Tipula carinifrons, 258
impacts of sea level rise, 636	Tlingit, 656, 681
projected impacts, 588	Tofieldia pusilla, 549
terrestrial ecosystems	Tomentypnum nitens, 257, 258
biodiversity monitoring, 586	Tonicella sp., 483
carbon storage, 305, 321	Toolik Lake, Alaska, 378, 390–391
characteristics, 1017	Tortula ruralis, 258
described, 11	tourism, 957, 963
feedback to climate change, 321–324, 550	and biodiversity, 579, 590
genetic diversity, 572–573	and boreal forests, 792, 794, 796
impacts of projected change in snow cover, 200	economic importance, 4, 15, 417, 675, 741–742, 963
impacts of projected permafrost change, 215	impacts of projected change in sea ice, 195
impacts of projected precipitation change, 188	impacts on Arctic ecosystems, 605, 639–640
impacts of trace gases, 312	impacts on indigenous peoples, 559, 657
impacts of UV radiation, 996	impacts on wildlife, 610, 616
and non-native species, 576	importance of wildlife, 600
paleoclimatic change, 249–253	and invasive species, 303
protected areas, 577	projected impacts, 195, 567, 1004, 1010, 1013
and surprises, 331	in protected areas, 604
trophic interactions, 298–304	role in mixed economies, 656
trophic structure, 549	toxaphene, 565, 891
uncertainties in projecting change, 329–334	trace gases, 312. see also carbon dioxide; methane; nitrous oxide
Teslin, Yukon Territory, 654	Trachurus trachurus. see horse mackerel
Tetrao urogallus. see wood grouse	traditional knowledge. see indigenous knowledge
Tetrastes bonasia. see hazel grouse	Trans Alaska Pipeline System, 932
Thaleichthys pacificus. see eulachon	transects, 821–824, 828–830
Thamnolia subuliformis, 258	transient response, 104
thaw frequency, 42–43	Transpolar Drift, 467, 471, 473
thaw settlement, 909, 914-916, 921, 927, 930-931, 932	transportation. see also navigation; Northern Sea Route; Northwest
Themisto libellula, 480, 482	Passage; winter roads
Theragra chalcogramma. see walleye pollock	and contaminants, 892
thermohaline circulation, 32–33, 54, 125, 461, 462	and habitat fragmentation, 609
and carbon cycle, 516	and human health, 866, 889
freshwater transport, 460	impacts of mild winter days, 23, 32
and glaciation, 47	impacts of projected change in river and lake ice, 223, 229
impacts of projected change, 473, 478	impacts of projected change in sea ice extent, 922–923
impacts of projected change in freshwater runoff, 229	impacts of projected change in snow cover, 200
and increased freshwater export, 223	impacts on arctic ecosystems, 605, 879, 880
and marine Arctic, 459	impacts on marine biodiversity, 576
observed change, 33, 45	impacts on wildlife, 616
paleoclimatic change, 33, 48, 50, 250, 1018	and natural hazards, 917
and past climate change, 125	and natural resource development, 937
projected change, 33, 119, 122, 124–125, 477–478, 519, 994,	observed damages to infrastructure, 936–937
999–1000, 1018	projected impacts, 566, 793–794, 920, 924–926, 927–928,
and sea ice, 196	930–931, 934–935, 948, 1000, 1001, 1002, 1007,
uncertainties, 521	1008, 1013
thermohaline ventilation, 458, 461–463, 473, 477–478	projected impacts and agriculture, 813
thermokarst, 927, 940, 1009. see also permafrost, terrestrial	tree rings. <i>see</i> paleoclimate; proxy climate records
and ecosystems, 245, 293	treeline
erosion, impacts on suspended sediment, 377	advance and albedo, 844
observed changes, 659	
and permafrost degradation, 210, 215, 319, 916	advance and tundra, 247, 550 and the Arctic Front, 319
projected impacts, 218, 254, 318, 435, 1012	barriers to movement, 999
thick-billed murre ( <i>Uria lomvia</i> ), 480, 489, 572, 623, 712	dynamics, 319–321
harvest by indigenous peoples, 653	historic change, 48, 51–52, 246–247, 368–371, 817, 846–851
harvest of, 626, 627, 629–630, 633	impacts of industrial forestry, 679–680
in northwest Atlantic, 732	impacts of overgrazing, 658
observed change, 1009	impacts of projected change on freshwater ecosystems, 370

impacts of projected change on indigenous peoples, 651	and altitude, 160, 163
indigenous observations of change, 993	and carbon balance, 310
and July temperature, 287	and cloud cover, 160, 162, 278
observed change, 1005, 1012	defined, 152–153
paleoclimatic, 251, 252, 253	dose rate, 153
and paludification, 550, 835	erythemal response spectrum, 153, 161–162
projected impacts, 253, 320–321, 570, 826, 851, 990, 991, 998,	factors affecting, 153, 159–165, 160
1006, 1012	and forage quality, 299
recent change in, 320, 847, 849-850	health impacts, 152–153, 174–175
trichinosis, 867, 880, 881	and herbivory, 853
Tringa erythropus. see spotted redshank	and human health, 874–878, 1000, 1001–1002
trout (Salmo spp.), 71, 695. see also individual species	impacts of increases, 996
trout perch (Percopsis omiscomaycus), 397	impacts of projected change, 998
Truncatella truncata, 298	impacts of projected change on animals, 281–282
Tryngites subruficollis. see buff-breasted sandpiper	impacts of projected change on biodiversity, 1017
tufted alpine saxifrage (Saxifraga caespitosa), 247	impacts of projected change on marine ecosystems, 512–516
tufted puffin (Fratercula cirrhata), 626, 633, 635	impacts of projected change on plants, 276–277
Tuktu and Nogak Project, 960–961	impacts on amphibians, 422
tularemia, 880, 881	impacts on aquatic ecosystems, 947, 999
tuna, 721. <i>see also</i> bluefin tuna	impacts on biogeochemical cycling, 513
tundra bean goose (Anser fabalis rossicus), 288, 289, 574	impacts on biota, 174–175
- '	*
tundra ecosystems, 294	impacts on boreal forests, 851–853
albedo, 550, 785	impacts on construction materials, 928–929
anthropogenic tundra, 320	impacts on crop plants, 852
biodiversity conservation, 548–551	impacts on fish, 413–415
carbon balance, 307	impacts on freshwater ecosystems, 419–427
carbon storage, 305	impacts on humans, 63, 419–420, 953
characteristics of, 244–247, 292–293	impacts on insect diseases, 302
classification of, 293	impacts on marine ecosystems, 520, 521–522
distribution in Russia, 790	impacts on materials, 175
extent and carbon balance, 322	impacts on photochemical reactions, 5
impacts of reindeer overgrazing, 657	impacts on plants, 571, 953
importance of lichens, 563	impacts on wildlife, 606, 953
and methane emissions, 318	indicators of, 899
methane fluxes, 312–313	indigenous observations, 63
net primary productivity, 306	indigenous observations of change, 168, 876
paleoclimatic, 252, 253	interaction with pollutants, 953
plant biomass, 306	irradiance, 153
projected change in area, 321, 569–570, 851, 990, 1008	and litter decomposition, 852–853
projected change in distribution, 999, 1005–1006	measurements, ground-based, 165–167
projected impacts, 253, 254, 327–328, 998, 1004, 1012	and mercury depletion events, 891
soil characteristics, 255	and moth defoliation of birch forests, 299
soil organic matter, 306	observed trends, 165–169, 248, 996
and treeline advance, 247, 550	and ozone depletion, 153, 154, 875, 995
trophic structure, 549	and ozone levels, 158, 160
tundra swan (Cygnus columbianus), 363, 617, 669	and pollutants, 877–878
tundra-taiga ecotone, 244, 319–321, 551–552, 788, 844	prehistoric, 253
biodiversity, 1016	projected change, 5–6, 174–175, 248, 253, 326, 419–420, 996
· · · · · · · · · · · · · · · · · · ·	
carbon storage, 305	radiation amplification factors, 161–162
characteristics, 292	and receptor orientation, 160, 164–165
impacts of projected change on indigenous peoples, 651	research recommendations, 175–176, 854
projected change in location, 569–570	satellite estimates, 168–169
recent change in, 849–850	and snow and ice cover, 160, 164, 998
in Russia, 789–790	and solar activity, 160, 160–161
in Sweden, 796	and solar zenith angle, 160, 161
wildfires in, 620	underwater exposure, 419–420
tundra vole (Microtus oeconomus), 258, 279, 280	and water quality, 160, 164
tundra-steppe. see steppe-tundra	umingmak. see muskox
turbellarians, 480	uncertainty. see also likelihood
turbot (Psetta maxima), 500, 716	about impacts of non-native species, 545
Turdus iliacus. see redwing	about impacts on caribou, 645
Turdus migratorius. see American robin	about terrestrial water balance, 31
Turdus pilaris. see fieldfare	in boreal forest role in carbon balance, 786
Tursiops truncatus. see bottlenose dolphin	in carbon balance projections, 999, 1015
tusk (Brosme brosme), 500, 711, 717	in climate simulations, 100–102, 114, 120, 121, 131, 140,
typhoid, 884	143–144, 477
••	in cloud simulations, 110, 118, 141
U	in estimating past UV radiation levels, 167
ultraviolet (UV) radiation. see also UV index	and forest management, 792–793, 794
action spectrum, 153, 875–876	in Greenland ice sheet mass budget, 205
and aerosols, 160, 162–163	in impacts on thermohaline circulation, 477–478
and albedo, 160, 163–164, 278	and infrastructure design, 938

methods for reducing, 10	indigenous observations of change, 272, 668
in observational data, 111, 112, 185, 197, 203, 226	observed change, 272
in ozone projections, 170, 173	present-day, 787
in paleoclimatic reconstructions, 47	projected change, 294, 322, 327–329, 568–570, 569, 569–570,
in projecting impacts, 2, 519	<i>588–589</i> , 605, <i>787</i> , 1017
in projecting impacts on fisheries, 693–694, 706–707,	projected change, approach, 323
739–740, 739–741, 770	in Russia, 790
in projecting impacts on freshwater ecosystems, 354–356, 385,	velvet scoter (Melanitta fusca), 86, 630
393, 394, 403–404, 434	vendace (Coregonus albula), 396
in projecting impacts on marine ecosystems, 507, 510	Vertagopus brevicaudus, 258, 259
	~ ·
in projecting impacts on terrestrial ecosystems, 329–334	Vitamin D, 877
in projections of sea level rise, 234	Voices from the Bay, 960
recommendations for reducing, 175, 188–189, 196, 208, 236,	voles (Microtus and Clethrionomys spp.), 549. see also individual
332–335, 520, 522	species
in snow cover simulations, 107	adaptation to arctic environment, 267–268
in UV projections, 174	echinococcus, 880
ungulates, 549, 570, 609, 1017. see also caribou; moose; reindeer	observed change in population cycles, 279–280, 301
United Nations Fish Stocks Agreement, 694–695	parasites and climate change, 302
United Nations Framework Convention on Climate Change, 119,	population cycles, 268, 281
956	and predation, 302
United Nations Law of the Sea Convention, 694	projected change, 998, 1006
Uria aalge. see common murre	projected impacts, 281
Uria lomvia, see thick-billed murre	trophic interactions, 298, 299–300, 301–302
Ursus americanus, see black bear	
	vulnerability
Ursus arctos. see brown bear	of arctic residents, 15, 195, 235–236, 865
Ursus maritimus. see polar bear	definition, 707, 947, 951
Urticina eques, 483	of indigenous peoples, 91–93, 664, 865, 866
Utsjoki. see Ochejohka, Finland	research recommendations, 95, 902, 1019–1020
UV index, 876. see also ultraviolet (UV) radiation	vulnerability assessments, 9
defined, 153	approaches, 949–957, 970–971, 980–981
increases in, 167	case studies, 961–980
maximum observed, 166	of human-environment systems, 946–951
,	methods, 957–961
V	and multiple stresses, 963
Vaccinium myrtillus. see blueberry	research recommendations, 982–983, 1019
Vaccinium spp., 293	Vulpes vulpes. see red fox
Vaccinium uliginosum. see bog whortleberry	
	Vuntut Gwitchin
Vaccinium vitis-idaea. see lingonberry	impacts case study, 680–685
Vanellus vanellus. see northern lapwing	
vegetation. see also plants; terrestrial ecosystems	W
adaptations to arctic climate, 557	waders, 490. see also birds, aquatic; seabirds
biomass, 306	biodiversity monitoring, 586
change and climate models, 107	diversity of, 549
change in distribution and POPs, 430	impacts of human activities, 564
growth forms, 247	migration routes, 556, 557, 564
impacts of ice crusting, 280	projected impacts, 574–575, 588, 604
impacts of projected change in permafrost, 215, 913	walleye (Sander vitreus), 395, 398, 405, 408, 411
impacts of projected change in snow cover, 200	walleye pollock (Theragra chalcogramma), 486, 495, 497
impacts of projected change in show cover, 200 impacts of projected change on indigenous peoples, 659	in Bering Sea, 748
impacts of projected change on microorganisms, 303	commercial fisheries, 652, 747, 748, 749–750, 762, 765,
impacts of projected change on regional climate, 314	766–767
impacts of projected change on water birds, 289, 290–291	distribution, 484
indigenous observations of change, 67, 271, 298, 660, 662, 685	harvest by indigenous peoples, 652
influence on water and energy exchange, 11	historic harvesting, 762
observed change, 294	impacts of past climate variability, 503, 503, 755–756, 766
observed change in biomass, 298	management regimes, 747, 763, 765
observed change in greenness, 298, 585	observed change, 1002
and permafrost, 909–910	as prey, 487
prehistoric change, 252	projected change in distribution, 545
prehistoric distribution, 251	projected impacts, 504, 507–508, 758, 759–760, 768, 1010
present-day, 4, 5, 245–246	temperature and distribution, 496
projected impacts, 246	,
1 3 1 /	walrus (Odobenus rosmarus), 487–488, 623–624
role in energy balance, 314	and benthos, 483
role in water balance, 314	in Bering Sea, 752
vegetation distribution	commercial harvest, 715, 753
and albedo, 322, 324	harvest by indigenous peoples, 652, 710
barriers to movement, 550, 568, 851	harvest of, 625, 627, 628, 632
and disturbance, 844	historic harvesting, 699, 761
and energy balance, 324	and human health, 882
feedbacks to climate system, 319–321	impacts of projected change in sea ice, 195, 659
historic change, 846–851	importance of sea ice, 632
impacts of projected change on reindeer/caribou, 660–661	indigenous observations of change, 70–71, 634

management agreements, 632	in northeast Atlantic, 696
migration, 546	in northwest Atlantic, 732
overharvesting, 487, 560, 1016	overharvesting, 487, 560, 1016
projected impacts, 510, 660, 999, 1010, 1011	productivity, 484
protected areas, 602	projected impacts, 380, 510–511, 520, 588, 660, 694, 999,
and sea ice, 497	1005, 1010
and trichinosis, 881	role in marine biodiversity, 561
on Wrangel Island, 604	threats from hunting, 614
water balance, terrestrial, 31–32	and tourism, 741
of boreal forest, 789	wheat, 807, 810, 852
of terrestrial ecosystems, 314, 318–319	whiskered auklet, 635
and vegetation distribution, 324	whistling swan. see tundra swan
water budgets	white-beaked dolphin (Lagenorhynchus albirostris), 489
of freshwater ecosystems, 357–358	white fox. see Arctic fox
water supply	white-fronted goose (Anser albifrons), 288, 290, 574, 669
impacts of projected change in glaciers and ice sheets, 208	white-rumped sandpiper (Calidris fuscicollis), 361
impacts of projected change in snow cover, 200	White Sea, 493
water vapor	white-sided dolphin (Lagenorhynchus acutus), 629
and hydrogen oxide (hydroxyl) radical, 155	white spruce (Picea glauca), 292
impacts of projected change in sea ice, 195	in Canada, 792
and ozone, 155	eastern spruce budworm outbreaks, 833
	*
stratospheric concentrations, 171	growth responses, 551–552
and stratospheric temperature, 155	projected change, 567, 824, 825–826, 828, 833
waterfowl. see birds, aquatic	reproduction and wildfire, 789
waterlogging. see paludification	spruce bark beetle outbreaks, 831–833
weasel, 258, 301	temperature impacts, 824, 825–826
weather. see also atmospheric pressure; extreme events;	tree ring data, 818–819
meteorological stations; precipitation; storms;	white-throated sparrow, 71
temperature, air	white wagtail (Motacilla alba), 258
impacts of projected change on subsistence activities, 667, 883	white whale. see beluga whale
indigenous observations of change, 68–69, 660, 661–662, 685,	white-winged scoter. see velvet scoter
869	whitefish (Coregonus spp.), 399, 406, 702, 704, 716, 762. see also
West Nile virus, 880, 881, 1011, 1014	cisco; cod; haddock; saithe
West Spitsbergen, 493	distribution, 396–397
Western Arctic Caribou Herd Working Group, 618–620	as fox prey, 76
western sandpiper (Calidris mauri), 288, 361	harvest by indigenous peoples, 652, 654
wetlands. see also freshwater ecosystems; peatlands	indigenous observations of change, 70
area of, 12	late freeze-up and harvests, 74–75
carbon balance, 364–365	paleoecology, 251
definition, 546	projected impacts, 373, 436
drainage impacts on biodiversity, 562	river preferences, 359
extent and methane emissions, 323	Whitehorse, Yukon Territory, 654
freshwater ecosystems of, 363–365	whooping crane (Grus americana), 418
impacts of changing UV radiation, 427	whortle-leaved willow (Salix myrsinites), 563
impacts of projected change on aquatic birds, 418	wildfires
impacts of projected change on biodiversity, 588	and biodiversity, 552–553
1 1 3 0	
impacts of projected precipitation/evaporation change, 188	in boreal forests, 785, 787, 789, 835–841
impacts of projected sea level rise, 235	and carbon balance, 841–843
and mercury production, 433	and caribou forage, 620, 647
methane emissions, 317, 999	and contaminants, 432
and permafrost, 45–46	in Fennoscandia, 839
projected change in area, 418	impacts of projected change in precipitation, 188
projected impacts, 381–384, 659, 998, 998, 1006, 1008, 1009	impacts on indigenous peoples, 678–680
projected impacts and aquatic birds, 617	increases in Russia, 608
projected impacts and addate birds, 617 projected impacts on carbon balance, 391–393, 434–435	management in Alaska, 838–839
	· · · · · · · · · · · · · · · · · · ·
sensitivity to permafrost degradation, 382	management in Canada, 837–838
whales, 489. see also individual species	management in Russia, 836–837
adaptation to cold, 556	observed change, 835, 837, 838–839, 1009
and benthos, 483	and permafrost thaw, 917
in Central North Atlantic, 711–712	projected impacts, 553, 838, 840–841, 999, 1007, 1008, 1012
food requirement, 484	and temperature, 803–804
harvest of, 627, 628, 629, 952	and weather and climate, 835–836
historic harvesting, 699, 701, 710, 715, 761, 766	wildlife. see also birds; mammals, marine; mammals, terrestrial
historic ranges, 624	corridors, 577–578
impacts of projected change in sea ice, 195, 659	defined, 543, 599
importance to Inuit, 651	economic importance, 1003
importance to tourism, 600	harvest in Russia, 609
indigenous harvesting, 710	and human settlement, 606
indigenous observations of change, 70–71	ice layers and grazing, 200
management agreements, 623, 633	impacts of harvesting, 601, 608, 610
management and indigenous subsistence activities, 665	impacts of industrial development, 620–621
management challenges, 601–602	impacts of tourism, 639–640

impacts of UV radiation, 953	woodpeckers, 291, 552
importance to tourism, 600	woolly rhinoceros (Coelodonta antiquitatis), 250
as indicators, 900	World Conservation Organization (IUCN), 575, 581–582, 585,
multiple stresses, 615	600, 603
population status in Alaska, 619 projected impacts, 599	Polar Bear Specialist Group, 623, 624, 625 World Meteorological Organization (WMO), 6
wildlife management	World Ocean Circulation Experiment (WOCE), 473
in Alaskan Arctic, 617–621, 632–634	worms, 540, 580
approaches, 601–602, 636–638	Wrangel Island Reserve, 604
in Canadian Arctic, 610–615, 626–628	
and climate change, 600, 604–605, 636–640	X
co-management, 612–613, 632–633, 636–637, 665–666,	Xanthoria candelaria, 269
670–671, 685, 960, 963, 1012	Xanthoria parietina, 270 Xema sabini. see Sabine's gull
co-management case study, 644–648 definition, 599	Xiphias gladius. see swordfish
in Fennoscandia, 616–617, 628–631	Alphus giunus. see swordiisii
future strategies, 634–636	Y
impacts on reindeer husbandry, 977	Yakutia, Russia, 609, 610-611
and indigenous subsistence activities, 664-665	Yamal Nenets, 962
and industrial development, 607	Yamal-Nenets Autonomous Okrug, 559, 675–678
and land and water use planning, 622, 638–639	Yamal Peninsula, 247, 256, 608
monitoring needs, 617	reindeer overgrazing, 657
present practices, 600–602 recommendations, 639–640	treeline change, 847–849 yellow-billed loon (Gavia adamsii), 363, 633, 635
research recommendations, 639–640	yellow-browed warbler ( <i>Phylloscopus inornatus</i> ), 258
role of indigenous peoples, 651	yellow Irish lord (Hemilepidotus jordani), 496, 756
role of protected areas, 602–605	"yellow legs." <i>see</i> white-fronted goose
in Russian Arctic, 606–610, 623–626, 638	yellow perch (Perca flavescens), 385-386, 395-396, 398, 404, 405
willow (Salix spp.), 257, 262, 293, 293-294, 303, 562. see also	yellowfin sole (Limanda aspera), 487
individual species	commercial fishery, 751
diversity of, 549	distribution, 484
in Greenland, 799	historic harvesting, 762
harvest by indigenous peoples, 654 in Iceland, 799–800	impacts of past climate variability, 756 Yenisey Basin, 185–188, 227
impacts of overgrazing, 563	Yenisey River
indigenous observations of change, 81, 271, 685	contaminant loading of, 381
willow grouse (Lagopus lagopus), 267	fish diversity in, 406–407
wind	freshwater discharge, 547
impacts on ocean mixing, 460–461, 544	freshwater ecosystems of, 358
impacts on species ranges, 588	impacts of impoundment, 358–359
indigenous observations of change, 68–69, 94, 632, 660, 661, 667, 684	observed trend in freeze-up date, 222 projected change in discharge, 228, 229
and infrastructure, 917	projected change in discharge, 226, 229 projected impacts, 358–359, 381
projected change, 470	ship traffic and wildlife, 609
and wildfires, 835	Yukaghir, 652
winter moth (Operophtera brumata), 299, 834	Yukon Basin, 227
winter roads. see also transportation	Yukon First Nations
impacts of projected change in river and lake ice, 223, 229	diet of indigenous peoples, 654, 657
observed change, 934	indigenous knowledge case study, 77–78
projected impacts, 793–794, 927, 933–934, 1001, 1004, 1010, 1013	subsistence activities, 653 Yukon River
witch (Glyptocephalus cynoglossus)	and carbon transport, 518
past distribution change, 500	freshwater ecosystems, 358
wolf (Canis lupus), 258, 259, 549	observed trend in breakup date, 222
harvest by indigenous peoples, 653, 667	projected change in discharge, 229
historic changes, 661	Yukon Territory, Canada
impacts of reindeer husbandry, 601	agriculture, 807–808
indigenous observations of change, 80	demographics, 868
in Norway and Sweden, 616 observed change, 1005	dietary contaminants, 657 indigenous observations of change, 68, 77–78
paleoecology, 252, 1016	indigenous subsistence production, 653
population monitoring, 620	Yupiit, 680
population status in Alaska, 619	Yup'ik, 652, 681
and reindeer/caribou, 301	
and reindeer/caribou predator control, 609, 615	Z
wolffish (Anarhichas lupus lupus), 711	zander (Sander lucioperca), 406, 407
wolverine (Gulo gulo), 258, 301, 549	zebra mussel ( <i>Dreissena polymorpha</i> ), 431
impacts of reindeer husbandry, 601, 609	zoobenthos, 479, 544
population status in Alaska, 619 as reindeer predator, 977	zooplankton, 711 adaptation to cold, 481
wood frog (Rana sylvatica), 422	in Bering Sea, 754
wood grouse (Tetrao urogallus), 86, 607	impacts of past climate variability, 497, 717

impacts of UV radiation, 513, 514
marine biomass, 479
in marine ecosystems, 482
and marine mammal reproduction, 496
marine productivity, 479
and ocean temperature, 755
projected impacts, 504, 506–507, 520, 544, 570
and sea ice extent, 505
secondary production, 493–494
and timing of phytoplankton bloom, 495